



Horikoshi, M., Beaumont, R., Day, F., Warrington, N., Kooijman, M., Ring, S. M., Hemani, G., McMahon, G., Sorensen, T., Lawlor, D. A., Davey Smith, G., Timpson, N. J., Evans, D., & Freathy, R. M., & Paternoster, L. (2016). Genome-wide associations for birth weight and correlations with adult disease. *Nature*, 538(7624), 248-252.
<https://doi.org/10.1038/nature19806>

Peer reviewed version

Link to published version (if available):
[10.1038/nature19806](https://doi.org/10.1038/nature19806)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Nature Publishing Group at doi:10.1038/nature19806. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Contents of Supplementary Information

1. Supplementary Tables and Notes

These tables and notes are provided in this document.

| | | |
|--------------------------|---|----|
| • Supplementary Table 1 | Description of studies contributing to trans-ancestry meta-analysis: ancestry group and country of origin, sample size, data collection methods, and birth weight summaries and exclusions. | 3 |
| • Supplementary Table 2 | Description of studies contributing to trans-ancestry meta-analysis: genotyping, quality control, pre-phasing, imputation, and association analysis. | 6 |
| • Supplementary Table 3 | BW association summary statistics for each component of the trans-ancestry meta-analysis of 153,781 individuals for lead SNPs at loci attaining genome-wide significance ($P < 5 \times 10^{-8}$). | 9 |
| • Supplementary Table 4 | Summary statistics for lead SNPs attaining genome-wide significant evidence ($P < 5 \times 10^{-8}$) of association with BW in European ancestry meta-analysis of up to 143,677 individuals and/or trans-ancestry meta-analysis of up to 153,781 individuals. | 18 |
| • Supplementary Table 5 | Loci with multiple distinct association signals attaining genome-wide significance ($P < 5 \times 10^{-8}$) in approximate conditional meta-analysis of 143,677 individuals of European ancestry, using 5,000 white British participants from UK BioBank as a reference for linkage disequilibrium. | 23 |
| • Supplementary Table 6 | Candidate gene(s) at birth weight loci. | 24 |
| • Supplementary Table 7 | Summary of 99% credible sets at 62 distinct autosomal association signals. | 28 |
| • Supplementary Table 8 | BW credible set enrichment for DNaseI hypersensitive sites. | 30 |
| • Supplementary Table 9 | Genetic variance explained. | 33 |
| • Supplementary Table 10 | Look-up of maternal genotype effect on offspring BW (unadjusted for foetal genotype) in up to 68,254 mothers for the 60 BW loci detected in foetal birth weight GWAS. | 34 |
| • Supplementary Table 11 | Maternal and foetal conditional analyses in 12,909 ALSPAC mother-child pairs. | 36 |
| • Supplementary Table 12 | Summary results of LD Score regression analyses between birth weight and various diseases, metabolic and anthropometric traits. | 38 |

| | | |
|--|---|----|
| • Supplementary Table 13 | Estimating the proportion of the BW-adult phenotype covariance attributable to genotyped SNPs in UK Biobank and the Northern Finland Birth Cohort (NFBC) 1966 data. | 42 |
| • Supplementary Table 14 | List of 77 reported imprinted regions used in the current analysis. | 45 |
| • Supplementary Table 15 | Comparison of variance in BW between individuals heterozygous and homozygous for each of the 59 autosomal BW index SNPs (plus <i>DLK1</i>) in 57,715 UK Biobank samples. | 47 |
| • Supplementary Table 16 | Parent-of-origin specific analysis at 59 autosomal BW loci (plus <i>DLK1</i>) in 4,908 ALSPAC mother-child pairs. | 49 |
| • Supplementary Table 18 | Effect of transmitted and untransmitted maternal haplotype scores on BW in 5,201 ALSPAC mother-child pairs. | 51 |
| • Supplementary Table 19 | Reciprocal approximate conditional analyses at <i>YKT6-GCK</i> variants associated with birth weight (BW) or fasting glucose (FG), in European ancestry meta-analysis of up to 143,677 individuals. | 52 |
| • Supplementary Figure 1 | BW association enrichment for DNaseI hypersensitive (DHS) sites in 128 cell types and four genic annotations. | 53 |
| • Supplementary Figure 2 | Estimates of phenotypic, genetic and residual correlations (a) and estimates of genetic and residual covariance (b) between birth weight and adult metabolic or anthropometric traits in UK Biobank ‘white-British’ samples (up to N=57,715). | 54 |
| • Full grants and funding supports for contributing studies and CHARGE Consortium Hematology Working Group | | 55 |
| • Grants and funding supports for individuals | | 59 |
| • Study acknowledgements | | 60 |

2. Supplementary Data

These table (excel spreadsheet) and plots (pdf file) are provided in separate documents.

- Supplementary Table 17 Association of BW signals with various adult metabolic and anthropometric traits. (GWAS look-ups)
- 60 regional plots for birth weight association

Supplementary Table 1. Description of studies contributing to trans-ancestry meta-analysis: ancestry group and country of origin, sample size, data collection methods, and birth weight summaries and exclusions.

(a) Component 1: European ancestry GWAS

| Study | Ancestry group | Country of origin | Year(s) of birth | Sample size (M/F) | Data collection | Phenotype exclusions | Mean (SD) birth weight (grams) | | | Median (IQR) GA (week) at delivery |
|---------------------------|----------------|-------------------|------------------|------------------------|---|---|--------------------------------|---------------|---------------|------------------------------------|
| | | | | | | | Males | Females | Combined | |
| 1958 British Birth Cohort | European | UK | 1958 | 4,595 (2,320/2,275) | Measured by midwives; supplemented with obstetric records and interviews with mothers | Multiple births, GA <37 weeks | 3439 (484) | 3277 (468) | 3359 (483) | 40 (39-41) |
| ALSPAC ^{a,b} | European | UK | ~1992 | 7,285 (3,722/3,563) | Identified from obstetric data, records from the ALSPAC measurers, and birth notification | Multiple births, GA <37 weeks, 5 SD winsorisation | 3553 (491) | 3423 (450) | 3490 (476) | 40 (40-41) |
| CHOP-Caucasian | European | USA | 1988-present | 9,405 (5,040/4,365) | Questionnaire and medical records | Multiple births, GA <37 weeks (when available) | 3447 (582) | 3343 (549) | 3398 (569) | N/A |
| CoLaus | European | Switzerland | 1928-1970 | 2,089 (892/1,197) | Self-reported as adults | N/A | 3490 (668) | 3250 (661) | 3352 (675) | N/A |
| COPSAC-2000 | European | Denmark | 1998-2001 | 352 (173/179) | Medical records | Multiple births, GA <37 weeks | N/A | N/A | 3555 (485) | 40 (39-41) |
| COPSAC-2010 | European | Denmark | 2008-2011 | 589 (306/283) | Medical records | Multiple births, GA <37 weeks | 3635 (483) | 3536 (474) | 3588 (481) | 40 (39-41) |
| COPSAC-REGISTRY | European | Denmark | 1987-1999 | 1,210 (804/406) | Medical Records | Multiple births, GA <37 weeks | 3609 (498) | 3443 (447) | 3553 (488) | 40 (39-41) |
| DNBC | European | Denmark | 1996-2003 | 915 (475/440) | Danish Medical Birth Register | Multiple births, GA <37 weeks, congenital abnormalities | 3767 (480) | 3625 (443) | 3699 (468) | 40 (40-41) |
| ERF | European | Netherlands | Various | 459 (187/272) | Interview | GA <37 weeks | 3161 (680) | 2955 (608) | 3039 (644) | N/A |
| EPIC | European | UK | 1993-1997 | 8,939 (3,448/5,491) | Self-reported | N/A | 3505 (786) | 3266 (750) | 3358 (772) | N/A |
| Fenland (GA+) | European | UK | 1950-1975 | 5,188 (2,088/3,100) | Self-reported as adults | GA described as "very pre-term" or "pre-term" | 3433 (638) | 3260 (594) | 3394 (555) | N/A |
| Fenland (GA-) | European | UK | 1950-1975 | 833 (509/324) | Self-reported as adults | None | 3465 (593) | 3154 (608) | 3354 (624) | N/A |
| Generation R | European | Netherlands | 2002-2006 | 2,701 (1,378/1,323) | Hospital records and community midwives | Multiple births, GA <37 weeks | 3628 (494) | 3518 (475) | 3574 (488) | 40 (39-41) |
| GINIplus & LISAplus (GA+) | European | Germany | 1996-1999 | 656 (360/296) | Parental report of medical records | Multiple births, GA <37 weeks, <2500g | 3498 (406) | 3376 (417) | 3443 (415) | 40 (39-41) |

| Study | Ancestry group | Country of origin | Year(s) of birth | Sample size (M/F) | Data collection | Phenotype exclusions | Mean (SD) birth weight (grams) | | | Median (IQR) GA (week) at delivery |
|----------------------------|----------------|-------------------|------------------|--|---|--|--------------------------------|---------------|----------------|------------------------------------|
| | | | | | | | Males | Females | Combined | |
| GINIplus & LISApplus (GA-) | European | Germany | 1996-1999 | 790 (391/399) | Parental report of medical records | Multiple births, GA <37 weeks, <2500g | 3499 (429) | 3348 (423) | 3423 (433) | N/A |
| GOYA | European | Denmark | 1943-1952 | 149/0 (obese), 141/0 (control) | School health records | N/A | N/A | N/A | 3553 (711) | N/A |
| HBCS | European | Finland | 1934-1944 | 1472 (639/833) | Birth records | Multiple births, GA <37 weeks | 3536 (460) | 3375 (439) | 3444 (454) | 40 (39-41) |
| INMA | European | Spain | 1997-2006 | 1,021 (527/494) | Well-trained midwives and nurses | None | 3362 (406) | 3188 (422) | 3278 (423) | 40 (39-41) |
| INTER99 | European | Denmark | 1939-1969 | 4,243 (1,981/2,262) | Measured by midwives and obtained from obstetric record registry | Multiple births, GA <37 weeks | 3505 (493) | 3370 (469) | 3433 (485) | N/A |
| Leipzig | European | Germany | 1985 - 2010 | 597 (304/293) | Questionnaire to mothers, documentation of medical screening examination if available | GA <37 weeks | 3573 (531) | 3480 (538) | 3527 (536) | 40 (39-40) |
| NEO | European | Netherlands | 1943-1963 | 504 (exact; 200/304), 3215 (range; 1,450/1,765) | Questionnaire | N/A | 3669 (1068) | 3236 (973) | 3514 (1271) | N/A |
| NFBC1966 | European | Finland | 1966 | 5,009 (2,393/2,616) | Measured in hospitals | Multiple births, GA <37 weeks or unknown | 3607 (506) | 3480 (466) | 3541 (489) | 40 (39-41) |
| NFBC1986 | European | Finland | 1986 | 4,680 (2,306/2,374) | Measured in hospitals | Multiple births, GA <37 weeks or unknown | 3626 (543) | 3519 (521) | 3572 (535) | 40 (39-40) |
| NTR | European | Netherlands | 1926-1998 | 1,265 (447/818) | Parental report or self-reported | Multiple births, GA <37 weeks | 3414 (619) | 3544 (630) | 3343 (601) | 40 (40-40) |
| ORCADES | European | Scotland | 1920-1991 | 960 (330/630) | Self-reported as adults | N/A | 3401 (607) | 3654 (685) | 3488 (640) | N/A |
| PANIC | European | Finland | 1999-2002 | 436 (231/205) | Medical records and parental questionnaire | Multiple births, GA <37 weeks | 3646 (488) | 3528 (444) | 3588 (474) | 40 (39-41) |
| RAINE | European | Australia | 1989-1991 | 1,347 (693/654) | Recorded at delivery by study personnel or obtained from hospital reports | Multiple births, GA <37 weeks | 3505 (471) | 3390 (462) | 3449 (470) | 40 (39-41) |
| SORBS | European | Germany | 1925-1988 | 298 (113/185) | Interview at recruitment | N/A | N/A | N/A | 3393 (673) | N/A |
| STRIP | European | Finland | 1989-1991 | 599 (311/288) | Medical records | Multiple births, GA <37 weeks | 3696 (471) | 3535 (443) | 3619 (465) | 40 (39-40) |
| TEENAGE (GA+) | European | Greece | 1993-1998 | 279 (126/153) | Measured by midwives or paediatricians; supplemented with data from mothers' interviews | GA <37 weeks | 3403 (467) | 3280 (421) | 3336 (445) | 40 (38-40) |
| TEENAGE (GA-) | European | Greece | 1993-1998 | 551 (234/317) | Measured by midwives or paediatricians; supplemented with data from mothers' interviews | N/A | 3398 (459) | 3298 (438) | 3341 (449) | N/A |
| TDCOB-cases | European | Denmark | 1987-2007 | 669 | Measured by midwives and registered in | Multiple births | 3682 | 3629 | 3660 | 40 (39-41) |

| | | | | | | | | | | |
|----------------|----------|---------|-----------|----------------------|--|---------------------------------------|---------------|---------------|---------------|------------|
| | | | | (391/278) | Danish Civil Registry | | (536) | (545) | (540) | |
| TDCOB-controls | European | Denmark | 1991-2006 | 560 (211/349) | Measured by midwives and registered in Danish Civil Registry | Multiple births | 3627 (517) | 3483 (485) | 3540 (502) | 40 (39-41) |
| YFS | European | Finland | 1962-1977 | 1,915 (861/1,054) | Mothers' interview | Multiple births, GA >3 weeks pre-term | 3648 (491) | 3510 (451) | 3572 (475) | N/A |

(b) Component 2: UK BioBank

| Study | Ancestry group | Country of origin | Year(s) of birth | Sample size (M/F) | Data collection | Phenotype exclusions | Mean (SD) birth weight (grams) | | | Median (IQR) GA (weeks) at delivery |
|------------|----------------|-------------------|------------------|---------------------------|-------------------------|--|--------------------------------|---------------|---------------|-------------------------------------|
| | | | | | | | Males | Females | Combined | |
| UK BioBank | European | UK | 2006-2010 | 67,786 (40,425/27,361) | Self-reported as adults | Multiple births, birth weight <2500g or >4000g | 3452 (416) | 3349 (417) | 3391 (420) | N/A |

(c) Component 3: Non-European ancestry GWAS

| Study | Ancestry group | Country of origin | Year(s) of birth | Sample size (M/F) | Data collection | Phenotype exclusions | Mean (SD) birth weight (grams) | | | Median (IQR) GA (weeks) at delivery |
|-------------------------|------------------|-------------------|------------------|------------------------|---|--|--------------------------------|---------------|---------------|-------------------------------------|
| | | | | | | | Males | Females | Combined | |
| CHOP-AA | African American | USA | 1988-present | 6,635 (3,343/3,292) | Questionnaire and medical records | Multiple births, GA <37 weeks (when available) | 3276 (554) | 3184 (535) | 3231 (546) | N/A |
| CLHNS | Filipino | Philippines | 1983-84 | 1,449 (755/694) | Local birth attendants | Multiple births, GA <37 weeks | 3067 (401) | 3018 (403) | 3043 (403) | 40 (38-40) |
| Generation R Turkish | Turkish | Netherlands | 2002-2006 | 420 (215/205) | Hospital records and community midwives | Multiple births, GA <37 weeks | 3477 (500) | 3369 (415) | 3424 (463) | 40 (39-41) |
| Generation R Moroccan | Moroccan | Netherlands | 2002-2006 | 365 (188/177) | Hospital records and community midwives | Multiple births, GA <37 weeks | 3642 (447) | 3417 (344) | 3533 (416) | 41 (40-41) |
| Generation R Surinamese | Surinamese | Netherlands | 2002-2006 | 395 (215/180) | Hospital records and community midwives | Multiple births, GA <37 weeks | 3288 (556) | 3130 (490) | 3216 (532) | 40 (39-41) |
| SCORM | Chinese | Singapore | 1992-1995 | 840 (420/420) | Documented medical record booklet | GA <37 weeks | 3229 (422) | 3182 (475) | 3205 (450) | 39 (38-40) |

M, Males; F, Females; GA, gestational age; IQR, interquartile range; N/A, not applicable; SD, standard deviation.

^aBoyd A, Golding J, Macleod J, Lawlor DA, Fraser A, et al. Cohort Profile: the 'children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* 42, 111-127 (2013).

^bThe study website contains details of all the data that is available through a fully searchable data dictionary (<http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary/>).

Supplementary Table 2. Description of studies contributing to trans-ancestry meta-analysis: genotyping, quality control, pre-phasing, imputation, and association analysis.

(a) Component 1: European ancestry GWAS

| Study | Genotyping array(s) ^a | Sample quality control | | SNP scaffold quality control | | | Prephasing software | Imputation | | Association analysis | | Lambda (M/F) |
|---------------------------|----------------------------------|------------------------|---|------------------------------|---------------------|-----------|---------------------|------------|-----------------|----------------------|--------------------------|--------------|
| | | Call rate | Additional filters | Call rate | HWE <i>P</i> -value | Frequency | | Software | Reference panel | Software | Covariates or adjustment | |
| 1958 British Birth Cohort | I550, I610 | None | Relatedness, ancestry outliers, sex discrepancy, identity, channel contrast | 95% | 1×10^{-4} | MAF<1% | MaCH | Minimac | 1000G Mar 2012 | ProbABEL | GA | 1.01/1.00 |
| ALSPAC | I550 | 97% | Heterozygosity, relatedness, ancestry outliers | 95% | 5×10^{-7} | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA, PC7 | 1.02/1.01 |
| CHOP-Caucasian | I550, I610 | 95% | Relatedness, ancestry outliers, sex discrepancy | 95% | 1×10^{-6} | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | PC1-3 | 0.96/0.96 |
| CoLaus | A5 | 90% | Relatedness, ancestry outliers | 90% | 1×10^{-7} | MAF<1% | MaCH | Minimac | 1000G Mar 2012 | In-house | None | 0.99/1.00 |
| COPSAC-2000 | I550 | 97.5% | Heterozygosity, relatedness, ancestry outliers | 98% | 1×10^{-6} | MAF<0.1% | MaCH | Minimac | 1000G Mar 2012 | mach2qtl | GA | 1.00/1.01 |
| COPSAC-2010 | IOEE | 95% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy | 95% | 1×10^{-6} | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | QuickTest | GA, PC1-5 | 1.01/1.00 |
| COPSAC-REGISTRY | IOEE | 95% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy | 97.5% | 1×10^{-6} | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | QuickTest | GA, PC1-5 | 1.00/1.00 |
| DNBC | I660 | 96% | Heterozygosity, ancestry outliers, sex discrepancy | 98% | 1×10^{-6} | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA | 1.00/1.00 |
| ERF | Various | 98% | Relatedness, ancestry outliers, sex discrepancy | 98% | 5×10^{-8} | MAF<0.5% | MaCH | Minimac | 1000G Mar 2012 | ProbABEL | Kinship matrix | 1.02/0.95 |
| EPIC | AUKBB | 97% | Heterozygosity, relatedness, sex discrepancy, singletons, channel contrast | 95% | 1×10^{-6} | MAC<1 | SHAPEIT | IMPUTE2 | 1000G Mar 2012 | SNPTEST | PC1-10 | 1.00/1.01 |
| Fenland (GA+) | AUKBB | 95% | Sex discrepancy, identity | 95% | 1×10^{-6} | MAC<2 | SHAPEIT | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA, PC1-10 | 1.00/1.01 |
| Fenland (GA-) | AUKBB | 95% | Sex discrepancy, identity | 95% | 1×10^{-6} | MAC<2 | SHAPEIT | IMPUTE2 | 1000G Mar 2012 | SNPTEST | PC1-10 | 0.99/1.00 |
| Generation R | I610, I660 | 97.5% | Heterozygosity, ancestry outliers, sex discrepancy | 98% | 1×10^{-6} | MAF<1% | MaCH | Minimac | 1000G Mar 2012 | mach2qtl | GA, PC1-4 | 1.03/1.01 |
| GINIplus & LISAPlus | A5, A6 | 95% | Heterozygosity, ancestry outliers, sex discrepancy | 95% | 1×10^{-5} | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA | 0.99/1.00 |
| GOYA | I610 | 95% | Heterozygosity, ancestry outliers, sex discrepancy | 95% | 1×10^{-7} | MAF<1% | MaCH | MaCH | 1000G Mar 2012 | QuickTest | None | 1.00/0.99 |
| HBCS | I670 | 95% | Heterozygosity, relatedness, ancestry outliers | 95% | 1×10^{-6} | MAF<1% | MaCH | MaCH | 1000G Mar 2012 | mach2qtl | GA | 1.02/1.02 |

| Study | Genotyping array(s) ^a | Sample quality control | | SNP scaffold quality control | | | Prephasing software | Imputation | | Association analysis | | Lambda (M/F) |
|---------------|----------------------------------|------------------------|--|------------------------------|--|-----------|---------------------|------------|-----------------|----------------------|--|--------------|
| | | Call rate | Additional filters | Call rate | HWE <i>p</i> -value | Frequency | | Software | Reference panel | Software | Covariates or adjustment | |
| INMA | IOQ | 98% | Heterozygosity, relatedness, ancestry outliers, duplicates | 95% | 1.1×10^{-6} | MAF<1% | IMPUTE2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA | 1.00/0.99 |
| INTER99 | ICM | 95% | Relatedness, ancestry outliers, sex discrepancy | 95% | 1×10^{-4} | MAF<1% | IMPUTE2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | PC1 | 0.97/1.02 |
| Leipzig | ICM | 95% | Duplicates, ancestry outliers, sex discrepancy | 95% (99% if MAF<5%) | 1×10^{-4} | MAF<1% | IMPUTE2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA | 0.95/0.96 |
| NEO | ICE | 98% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy | 98% | 1×10^{-6} | None | IMPUTE2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | PC1-5 | 0.99/0.99 |
| NFBC1966 | I370 | 95% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy, duplicates, withdrawn consent | 95% (99% if MAF<5%) | 5.7×10^{-7} | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA, PC1-3 | 1.00/0.99 |
| NFBC1986 | ICM | 95% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy, duplicates, withdrawn consent | 95% (99% if MAF<5%) | 5.7×10^{-7} (1×10^{-4} if MAF<5%) | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA, PC1-3 | 1.00/1.10 |
| NTR | A6, I370, I660, IOQ | 90% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy | 95% | 1×10^{-5} | MAF<1% | MaCH | Minimac | 1000G Mar 2012 | PLINK | GA, array, PC1-6 (global), PC1-3 (local) | 1.08/1.04 |
| ORCADES | I300, IOQ, IOE | 95% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy, duplicates | 95% (99% if MAF<5%) | 1×10^{-6} | MAF<1% | SHAPEIT | IMPUTE2 | 1000G Mar 2012 | ProbABEL | array, PC1-3 | 1.00/0.99 |
| PANIC | ICM, ICE | 90 % | Heterozygosity, relatedness, ancestry outliers, sex discrepancy | 95% | 1×10^{-6} | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA, PC1-4 | 1.01/1.01 |
| RAINE | I660 | 97% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy, chromosomal abnormalities | 95% | 5.7×10^{-7} | MAF<1% | MaCH | MaCH | 1000G Mar 2012 | ProbABEL | GA, PC1-2 | 1.01/0.99 |
| SORBS | I660 | 94% | Relatedness, ancestry outliers, sex discrepancy, duplicates | 95% | 1×10^{-4} | MAF<1% | MaCH | Minimac | 1000G Mar 2012 | ProbABEL | Kinship matrix | 1.01/1.01 |
| STRIP | A5, A6 | 95% | Heterozygosity, ancestry outliers, twins | 95% | 1×10^{-6} | MAF<0.1% | SHAPEIT | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA, PC1-4 | 1.01/1.01 |
| TEENAGE (GA+) | ICM | 95% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy | 95% (99% if MAF<5%) | 1×10^{-4} | MAF<1% | SHAPEIT | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA | 1.02/1.00 |
| TEENAGE (GA-) | IOE | 95% | Heterozygosity, relatedness, ancestry outliers, sex discrepancy | 95% (99% if MAF<5%) | 1×10^{-4} | MAF<1% | SHAPEIT | IMPUTE2 | 1000G Mar 2012 | SNPTEST | None | 1.02/0.98 |

| | | | | | | | | | | | | |
|----------------|------|-----|--|-----|--------------------|--------|----------|---------|----------------|---------|---------|-----------|
| TDCOB-cases | IOE | 95% | Heterozygosity, relatedness, ancestry outliers | 95% | 1x10 ⁻⁶ | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA, PC1 | 1.02/1.01 |
| TDCOB-controls | ICE | 95% | Heterozygosity, relatedness, ancestry outliers | 95% | 1x10 ⁻⁶ | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA, PC1 | 1.02/1.05 |
| YFS | I670 | 95% | Heterozygosity, relatedness, sex discrepancy, duplicates | 95% | 1x10 ⁻⁶ | MAF<1% | SHAPEIT | IMPUTE2 | 1000G Mar 2012 | SNPTEST | PC1-4 | 0.99/1.01 |

(b) Component 2: UK BioBank

| Study | Genotyping array(s) ^a | Sample quality control | | SNP scaffold quality control | | | Prephasing software | Imputation | | Association analysis | | Lambda |
|------------|----------------------------------|------------------------|--|------------------------------|---------------------|-----------|---------------------|------------|------------------------|----------------------|--------------------------|--------|
| | | Call rate | Additional filters | Call rate | HWE <i>P</i> -value | Frequency | | Software | Reference panel | Software | Covariates or adjustment | |
| UK BioBank | AUKBB | 98% | Heterozygosity, relatedness, ancestry outliers | 95% | N/A | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Oct 2014 & UK10K | BOLT-LMM | Sex, genotype array | N/A |

(c) Component 3: Non-European ancestry GWAS.

| Study | Genotyping array(s) ^a | Sample quality control | | SNP scaffold quality control | | | Prephasing software | Imputation | | Association analysis | | Lambda (M/F) |
|-------------------------|----------------------------------|------------------------|--|------------------------------|---------------------|-----------|---------------------|------------|-----------------|----------------------|--------------------------|--------------|
| | | Call rate | Additional filters | Call rate | HWE <i>P</i> -value | Frequency | | Software | Reference panel | Software | Covariates or adjustment | |
| CHOP-AA | I550, I610 | 95% | Relatedness, ancestry outliers, sex discrepancy | 95% | 1x10 ⁻⁶ | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | PC1-3 | 0.98/0.98 |
| CLHNS | ICM | 98.6% | Relatedness, sex discrepancy | 97% | 1x10 ⁻⁶ | N/A | MaCH | MaCH | 1000G Mar 2012 | mach2qtl | GA | 1.02/1.02 |
| Generation R Turkish | I610, I660 | 97.5% | Heterozygosity, ancestry outliers, sex discrepancy | 95% | 1x10 ⁻⁷ | MAF<1% | MaCH | Minimac | 1000G Mar 2012 | mach2qtl | GA, PC1-4 | 1.01/1.02 |
| Generation R Moroccan | I610, I660 | 97.5% | Heterozygosity, ancestry outliers, sex discrepancy | 90% | 1x10 ⁻⁷ | MAF<1% | MaCH | Minimac | 1000G Mar 2012 | mach2qtl | GA, PC1-4 | 1.01/0.98 |
| Generation R Surinamese | I610, I660 | 97.5% | Heterozygosity, ancestry outliers, sex discrepancy | 98% | 1x10 ⁻⁷ | MAF<1% | MaCH | Minimac | 1000G Mar 2012 | mach2qtl | GA, PC1 | 0.99/0.95 |
| SCORM | I550 | 95% | Heterozygosity, relatedness, sex discrepancy | 95% | 1x10 ⁻⁶ | MAF<1% | SHAPEIT2 | IMPUTE2 | 1000G Mar 2012 | SNPTEST | GA | 0.98/0.99 |

HWE, Hardy-Weinberg equilibrium; MAF, minor allele frequency; MAC, minor allele count; GA, gestational age; PC, principal component.

^aGenotype array codes: Affymetrix 5.0 (A5); Affymetrix 6.0 (A6); Affymetrix Axiom UK BiLEVE (AUKBL); Affymetrix Axiom UK BioBank (AUKBB); Illumina Human370CNV (I370); Illumina HumanHap550 (I550); Illumina HumanHap610 (I610); Illumina HumanHap660 (I660); Illumina HumanHap670 (I670); Illumina CardioMetaboChip (ICM); Illumina OmniQuad (IOQ); Illumina OmniExpress (IOE); Illumina CoreExome (ICE); Illumina OmniExpressExome (IOEE).

Supplementary Table 3. BW association summary statistics for each component of the trans-ancestry meta-analysis of 153,781 individuals for lead SNPs at loci attaining genome-wide significance ($P < 5 \times 10^{-8}$).

Locus: *WNT4-ZBTB40*. Lead SNP: rs2473248. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.86 | 0.039 | 0.008 | 5.0×10^{-7} | 71,642 | 0.66 |
| Component 2: UKBB | 0.88 | 0.025 | 0.008 | 0.0032 | 67,786 | N/A |
| European ancestry meta-analysis | 0.87 | 0.033 | 0.006 | 1.1×10^{-8} | 139,428 | 0.21 |
| Component 3: non-European ancestry | 0.78 | 0.039 | 0.018 | 0.034 | 8,653 | 0.43 |
| Trans-ancestry meta-analysis | 0.87 | 0.033 | 0.005 | 1.1×10^{-9} | 148,081 | 0.72 |

Locus: *ZBTB7B*. Lead SNP: rs3753639. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.23 | 0.038 | 0.006 | 3.9×10^{-9} | 70,376 | 0.60 |
| Component 2: UKBB | 0.25 | 0.024 | 0.006 | 0.00017 | 67,786 | N/A |
| European ancestry meta-analysis | 0.24 | 0.031 | 0.004 | 7.3×10^{-12} | 138,162 | 0.12 |
| Component 3: non-European ancestry | 0.18 | 0.037 | 0.020 | 0.061 | 8,655 | 0.40 |
| Trans-ancestry meta-analysis | 0.23 | 0.031 | 0.004 | 1.3×10^{-12} | 146,817 | 0.72 |

Locus: *FCGR2B*. Lead SNP: rs72480273. Effect/other alleles: C/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.16 | 0.029 | 0.007 | 8.7×10^{-5} | 70,594 | 0.14 |
| Component 2: UKBB | 0.19 | 0.033 | 0.007 | 2.3×10^{-6} | 67,786 | N/A |
| European ancestry meta-analysis | 0.17 | 0.031 | 0.005 | 8.0×10^{-10} | 138,380 | 0.69 |
| Component 3: non-European ancestry | 0.15 | 0.005 | 0.023 | 0.84 | 8,655 | 0.71 |
| Trans-ancestry meta-analysis | 0.17 | 0.030 | 0.005 | 1.5×10^{-9} | 147,035 | 0.86 |

Locus: *DTL*. Lead SNP: rs61830764. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.38 | 0.031 | 0.006 | 1.0×10^{-7} | 70,372 | 0.12 |
| Component 2: UKBB | 0.38 | 0.013 | 0.006 | 0.019 | 67,786 | N/A |
| European ancestry meta-analysis | 0.38 | 0.022 | 0.004 | 5.6×10^{-8} | 138,158 | 0.029 |
| Component 3: non-European ancestry | 0.15 | 0.016 | 0.024 | 0.50 | 8,655 | 0.30 |
| Trans-ancestry meta-analysis | 0.36 | 0.022 | 0.004 | 4.5×10^{-8} | 146,813 | 0.18 |

Locus: *ATAD2B*. Lead SNP: rs7575873. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.88 | 0.040 | 0.008 | 4.9×10^{-7} | 71,639 | 0.98 |
| Component 2: UKBB | 0.87 | 0.036 | 0.008 | 6.3×10^{-6} | 67,786 | N/A |
| European ancestry meta-analysis | 0.88 | 0.038 | 0.006 | 1.3×10^{-11} | 139,425 | 0.71 |
| Component 3: non-European ancestry | 0.93 | -0.011 | 0.027 | 0.67 | 10,104 | 0.61 |
| Trans-ancestry meta-analysis | 0.88 | 0.036 | 0.006 | 6.2×10^{-11} | 149,529 | 0.35 |

Locus: *EPAS1*. Lead SNP: rs1374204. Effect/other alleles: T/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.70 | 0.050 | 0.006 | 1.0×10^{-16} | 66,667 | 0.94 |
| Component 2: UKBB | 0.70 | 0.044 | 0.006 | 9.7×10^{-14} | 67,786 | N/A |
| European ancestry meta-analysis | 0.70 | 0.047 | 0.004 | 6.2×10^{-29} | 134,453 | 0.42 |
| Component 3: non-European ancestry | 0.63 | 0.031 | 0.016 | 0.049 | 8,654 | 0.14 |
| Trans-ancestry meta-analysis | 0.70 | 0.046 | 0.004 | 1.5×10^{-29} | 143,107 | 0.82 |

Locus: *PTH1R*. Lead SNP: rs2242116. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.38 | 0.018 | 0.005 | 0.00048 | 75,884 | 0.48 |
| Component 2: UKBB | 0.38 | 0.025 | 0.006 | 5.5×10^{-6} | 67,786 | N/A |
| European ancestry meta-analysis | 0.38 | 0.022 | 0.004 | 1.4×10^{-8} | 143,670 | 0.37 |
| Component 3: non-European ancestry | 0.57 | 0.012 | 0.014 | 0.41 | 10,103 | 0.92 |
| Trans-ancestry meta-analysis | 0.39 | 0.021 | 0.004 | 1.2×10^{-8} | 153,773 | 0.94 |

Locus: *ADCY5*. Lead SNP: rs11719201. Effect/other alleles: T/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.22 | 0.052 | 0.006 | 5.2×10^{-18} | 75,884 | 0.34 |
| Component 2: UKBB | 0.24 | 0.039 | 0.006 | 4.0×10^{-10} | 67,786 | N/A |
| European ancestry meta-analysis | 0.23 | 0.046 | 0.004 | 2.4×10^{-26} | 143,670 | 0.14 |
| Component 3: non-European ancestry | 0.13 | 0.037 | 0.022 | 0.095 | 9,264 | 0.18 |
| Trans-ancestry meta-analysis | 0.23 | 0.046 | 0.004 | 6.4×10^{-27} | 152,934 | 0.35 |

Locus: *CPA3*. Lead SNP: rs10935733. Effect/other alleles: T/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.42 | 0.023 | 0.005 | 2.2×10^{-5} | 71,640 | 0.58 |
| Component 2: UKBB | 0.39 | 0.021 | 0.005 | 0.00012 | 67,786 | N/A |
| European ancestry meta-analysis | 0.41 | 0.022 | 0.004 | 9.2×10^{-9} | 139,426 | 0.80 |
| Component 3: non-European ancestry | 0.60 | 0.038 | 0.015 | 0.013 | 10,103 | 0.62 |
| Trans-ancestry meta-analysis | 0.42 | 0.023 | 0.004 | 6.2×10^{-10} | 149,529 | 0.33 |

Locus: *CCNL1-LEKR1*. Lead SNP: rs13322435. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.60 | 0.063 | 0.005 | 1.0×10^{-30} | 71,640 | 0.0091 |
| Component 2: UKBB | 0.60 | 0.042 | 0.006 | 3.4×10^{-13} | 67,786 | N/A |
| European ancestry meta-analysis | 0.60 | 0.053 | 0.004 | 3.7×10^{-41} | 139,426 | 0.0080 |
| Component 3: non-European ancestry | 0.45 | 0.040 | 0.015 | 0.0066 | 10,103 | 0.66 |
| Trans-ancestry meta-analysis | 0.59 | 0.052 | 0.004 | 1.3×10^{-42} | 149,529 | 0.14 |

Locus: *LCORL*. Lead SNP: rs925098. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.28 | 0.040 | 0.006 | 4.8×10^{-12} | 71,640 | 0.46 |
| Component 2: UKBB | 0.26 | 0.027 | 0.006 | 9.0×10^{-6} | 67,786 | N/A |
| European ancestry meta-analysis | 0.27 | 0.034 | 0.004 | 5.4×10^{-16} | 139,426 | 0.11 |
| Component 3: non-European ancestry | 0.34 | 0.011 | 0.015 | 0.44 | 10,102 | 0.28 |
| Trans-ancestry meta-analysis | 0.28 | 0.032 | 0.004 | 1.3×10^{-15} | 149,528 | 0.060 |

Locus: *HHIP*. Lead SNP: rs6537307. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.50 | 0.023 | 0.005 | 1.5×10^{-5} | 71,645 | 0.35 |
| Component 2: UKBB | 0.49 | 0.029 | 0.005 | 1.1×10^{-7} | 67,786 | N/A |
| European ancestry meta-analysis | 0.50 | 0.025 | 0.004 | 9.5×10^{-12} | 139,431 | 0.41 |
| Component 3: non-European ancestry | 0.21 | 0.038 | 0.018 | 0.036 | 9,557 | 0.92 |
| Trans-ancestry meta-analysis | 0.48 | 0.026 | 0.004 | 1.3×10^{-12} | 148,988 | 0.80 |

Locus: 5q11.2. Lead SNP: rs854037. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.81 | 0.038 | 0.007 | 7.8×10^{-9} | 71,643 | 0.021 |
| Component 2: UKBB | 0.82 | 0.014 | 0.007 | 0.045 | 67,786 | N/A |
| European ancestry meta-analysis | 0.81 | 0.027 | 0.005 | 2.2×10^{-8} | 139,429 | 0.011 |
| Component 3: non-European ancestry | 0.53 | 0.009 | 0.014 | 0.52 | 10,103 | 0.29 |
| Trans-ancestry meta-analysis | 0.80 | 0.025 | 0.005 | 3.5×10^{-8} | 149,532 | 0.010 |

Locus: *EBF1*. Lead SNP: rs7729301. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.72 | 0.021 | 0.006 | 0.00035 | 75,883 | 0.23 |
| Component 2: UKBB | 0.74 | 0.027 | 0.006 | 1.0×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.73 | 0.024 | 0.004 | 1.6×10^{-8} | 143,669 | 0.48 |
| Component 3: non-European ancestry | 0.58 | 0.033 | 0.014 | 0.022 | 10,104 | 0.64 |
| Trans-ancestry meta-analysis | 0.72 | 0.025 | 0.004 | 1.3×10^{-9} | 153,773 | 0.78 |

Locus: *CDKAL1*. Lead SNP: rs35261542. Effect/other alleles: C/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.72 | 0.047 | 0.006 | 6.1×10^{-17} | 75,881 | 0.94 |
| Component 2: UKBB | 0.74 | 0.041 | 0.006 | 1.4×10^{-11} | 67,786 | N/A |
| European ancestry meta-analysis | 0.73 | 0.044 | 0.004 | 4.4×10^{-27} | 143,667 | 0.49 |
| Component 3: non-European ancestry | 0.75 | 0.044 | 0.016 | 0.0059 | 10,102 | 0.35 |
| Trans-ancestry meta-analysis | 0.73 | 0.044 | 0.004 | 9.7×10^{-29} | 153,769 | 0.42 |

Locus: *HIST1H2BE*. Lead SNP: rs9379832. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.67 | 0.029 | 0.006 | 7.2×10^{-7} | 70,375 | 1.0 |
| Component 2: UKBB | 0.73 | 0.016 | 0.006 | 0.011 | 67,786 | N/A |
| European ancestry meta-analysis | 0.70 | 0.023 | 0.004 | 6.6×10^{-8} | 138,161 | 0.13 |
| Component 3: non-European ancestry | 0.77 | 0.044 | 0.021 | 0.038 | 10,103 | 0.70 |
| Trans-ancestry meta-analysis | 0.71 | 0.024 | 0.004 | 1.2×10^{-8} | 148,264 | 0.42 |

Locus: *HMGA1*. Lead SNP: rs7742369. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.17 | 0.027 | 0.007 | 0.0011 | 69,259 | 0.90 |
| Component 2: UKBB | 0.18 | 0.030 | 0.007 | 2.4×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.18 | 0.028 | 0.005 | 1.0×10^{-8} | 137,045 | 0.77 |
| Component 3: non-European ancestry | 0.44 | 0.013 | 0.016 | 0.41 | 9,561 | 0.18 |
| Trans-ancestry meta-analysis | 0.19 | 0.027 | 0.005 | 1.1×10^{-8} | 146,606 | 0.16 |

Locus: *L3MBTL3*. Lead SNP: rs1415701. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.75 | 0.025 | 0.006 | 2.3×10^{-5} | 75,880 | 0.69 |
| Component 2: UKBB | 0.73 | 0.025 | 0.006 | 3.2×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.74 | 0.025 | 0.004 | 2.6×10^{-9} | 143,666 | 1.00 |
| Component 3: non-European ancestry | 0.63 | 0.048 | 0.015 | 0.0013 | 10,102 | 0.64 |
| Trans-ancestry meta-analysis | 0.73 | 0.027 | 0.004 | 4.0×10^{-11} | 153,768 | 0.40 |

Locus: *ESR1*. Lead SNP: rs1101081. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.72 | 0.044 | 0.006 | 5.4×10^{-14} | 71,641 | 0.95 |
| Component 2: UKBB | 0.73 | 0.031 | 0.006 | 2.7×10^{-7} | 67,786 | N/A |
| European ancestry meta-analysis | 0.72 | 0.038 | 0.004 | 1.6×10^{-19} | 139,427 | 0.14 |
| Component 3: non-European ancestry | 0.79 | 0.027 | 0.017 | 0.13 | 10,103 | 0.47 |
| Trans-ancestry meta-analysis | 0.73 | 0.037 | 0.004 | 6.1×10^{-20} | 149,530 | 0.28 |

Locus: *GNA12*. Lead SNP: rs798489. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.73 | 0.020 | 0.006 | 0.00040 | 75,884 | 0.75 |
| Component 2: UKBB | 0.73 | 0.027 | 0.006 | 1.0×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.73 | 0.023 | 0.004 | 2.0×10^{-8} | 143,670 | 0.44 |
| Component 3: non-European ancestry | 0.92 | 0.053 | 0.027 | 0.048 | 9,708 | 0.14 |
| Trans-ancestry meta-analysis | 0.74 | 0.024 | 0.004 | 5.0×10^{-9} | 153,378 | 0.84 |

Locus: *IGF2BP3*. Lead SNP: rs11765649. Effect/other alleles: T/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.76 | 0.021 | 0.006 | 0.00049 | 71,642 | 0.80 |
| Component 2: UKBB | 0.74 | 0.032 | 0.006 | 1.3×10^{-7} | 67,786 | N/A |
| European ancestry meta-analysis | 0.75 | 0.027 | 0.004 | 5.8×10^{-10} | 139,428 | 0.19 |
| Component 3: non-European ancestry | 0.87 | 0.003 | 0.022 | 0.88 | 10,103 | 0.37 |
| Trans-ancestry meta-analysis | 0.76 | 0.026 | 0.004 | 1.0×10^{-9} | 149,531 | 0.60 |

Locus: *TBX20*. Lead SNP: rs6959887. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.61 | 0.031 | 0.005 | 1.4×10^{-9} | 75,873 | 0.58 |
| Component 2: UKBB | 0.61 | 0.013 | 0.006 | 0.021 | 67,786 | N/A |
| European ancestry meta-analysis | 0.61 | 0.023 | 0.004 | 1.5×10^{-9} | 143,659 | 0.013 |
| Component 3: non-European ancestry | 0.63 | -0.007 | 0.014 | 0.65 | 10,098 | 0.62 |
| Trans-ancestry meta-analysis | 0.61 | 0.021 | 0.004 | 1.0×10^{-8} | 153,757 | 0.054 |

Locus: *YKT6-GCK*. Lead SNP: rs138715366. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|--------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.9908 | 0.227 | 0.034 | 1.7×10^{-11} | 64,557 | 0.86 |
| Component 2: UKBB | 0.9915 | 0.253 | 0.031 | 7.1×10^{-16} | 67,786 | N/A |
| European ancestry meta-analysis | 0.9911 | 0.241 | 0.023 | 7.2×10^{-26} | 132,343 | 0.57 |
| Component 3: non-European ancestry | 0.9977 | 0.855 | 0.333 | 0.010 | 3,292 | 1.00 |
| Trans-ancestry meta-analysis | 0.9913 | 0.244 | 0.023 | 1.4×10^{-26} | 135,635 | 0.16 |

Locus: *MLXIPL*. Lead SNP: rs62466330. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.06 | 0.052 | 0.011 | 1.9×10^{-6} | 74,414 | 0.24 |
| Component 2: UKBB | 0.07 | 0.046 | 0.011 | 1.5×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.07 | 0.049 | 0.008 | 1.2×10^{-10} | 142,200 | 0.70 |
| Component 3: non-European ancestry | 0.04 | 0.142 | 0.045 | 0.0015 | 7,815 | 0.52 |
| Trans-ancestry meta-analysis | 0.07 | 0.051 | 0.007 | 5.9×10^{-12} | 150,015 | 0.23 |

Locus: *ANK1-NKX6-3*. Lead SNP: rs13266210. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.78 | 0.023 | 0.006 | 0.00034 | 71,643 | 0.19 |
| Component 2: UKBB | 0.79 | 0.040 | 0.007 | 1.5×10^{-9} | 67,786 | N/A |
| European ancestry meta-analysis | 0.79 | 0.031 | 0.005 | 1.3×10^{-11} | 139,429 | 0.054 |
| Component 3: non-European ancestry | 0.81 | 0.013 | 0.019 | 0.48 | 9,562 | 0.30 |
| Trans-ancestry meta-analysis | 0.79 | 0.030 | 0.004 | 1.6×10^{-11} | 148,991 | 0.32 |

Locus: *TRIB1*. Lead SNP: rs6989280. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.73 | 0.026 | 0.006 | 5.6×10^{-6} | 75,885 | 0.34 |
| Component 2: UKBB | 0.73 | 0.017 | 0.006 | 0.0069 | 67,786 | N/A |
| European ancestry meta-analysis | 0.73 | 0.022 | 0.004 | 2.2×10^{-7} | 143,671 | 0.25 |
| Component 3: non-European ancestry | 0.32 | 0.027 | 0.016 | 0.085 | 10,104 | 0.85 |
| Trans-ancestry meta-analysis | 0.70 | 0.022 | 0.004 | 5.0×10^{-8} | 153,775 | 0.92 |

Locus: *SLC45A4*. Lead SNP: rs12543725. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.59 | 0.023 | 0.005 | 1.1×10^{-5} | 71,645 | 0.39 |
| Component 2: UKBB | 0.59 | 0.023 | 0.005 | 3.0×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.59 | 0.023 | 0.004 | 1.2×10^{-9} | 139,431 | 0.94 |
| Component 3: non-European ancestry | 0.78 | 0.005 | 0.018 | 0.78 | 8,653 | 0.42 |
| Trans-ancestry meta-analysis | 0.60 | 0.022 | 0.004 | 1.9×10^{-9} | 148,084 | 0.65 |

Locus: *PTCH1*. Lead SNP: rs28510415. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.09 | 0.041 | 0.009 | 1.4×10^{-5} | 66,960 | 0.40 |
| Component 2: UKBB | 0.10 | 0.070 | 0.009 | 1.7×10^{-14} | 67,786 | N/A |
| European ancestry meta-analysis | 0.09 | 0.056 | 0.007 | 1.5×10^{-17} | 134,746 | 0.023 |
| Component 3: non-European ancestry | 0.03 | -0.086 | 0.043 | 0.049 | 8,654 | 0.79 |
| Trans-ancestry meta-analysis | 0.09 | 0.053 | 0.006 | 4.0×10^{-16} | 143,400 | 0.0055 |

Locus: *LPAR1*. Lead SNP: rs2150052. Effect/other alleles: T/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.50 | 0.021 | 0.005 | 8.6×10^{-5} | 71,638 | 0.23 |
| Component 2: UKBB | 0.51 | 0.021 | 0.005 | 7.5×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.50 | 0.021 | 0.004 | 2.2×10^{-8} | 139,424 | 0.95 |
| Component 3: non-European ancestry | 0.46 | 0.008 | 0.015 | 0.61 | 8,655 | 0.71 |
| Trans-ancestry meta-analysis | 0.50 | 0.020 | 0.004 | 2.8×10^{-8} | 148,079 | 0.99 |

Locus: *PHF19*. Lead SNP: rs7847628. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.68 | 0.028 | 0.006 | 1.1×10^{-6} | 71,638 | 0.47 |
| Component 2: UKBB | 0.68 | 0.018 | 0.006 | 0.0014 | 67,786 | N/A |
| European ancestry meta-analysis | 0.68 | 0.023 | 0.004 | 1.0×10^{-8} | 139,424 | 0.23 |
| Component 3: non-European ancestry | 0.57 | 0.017 | 0.015 | 0.25 | 10,104 | 0.30 |
| Trans-ancestry meta-analysis | 0.67 | 0.023 | 0.004 | 5.4×10^{-9} | 149,528 | 0.17 |

Locus: *STRBP*. Lead SNP: rs700059. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.15 | 0.027 | 0.007 | 0.00028 | 71,641 | 0.090 |
| Component 2: UKBB | 0.14 | 0.040 | 0.008 | 2.1×10^{-7} | 67,786 | N/A |
| European ancestry meta-analysis | 0.14 | 0.033 | 0.005 | 4.7×10^{-10} | 139,427 | 0.22 |
| Component 3: non-European ancestry | 0.43 | 0.059 | 0.016 | 0.00018 | 8,653 | 0.34 |
| Trans-ancestry meta-analysis | 0.16 | 0.036 | 0.005 | 1.2×10^{-12} | 148,080 | 0.21 |

Locus: *HHEX-IDE*. Lead SNP: rs61862780. Effect/other alleles: T/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.51 | 0.027 | 0.005 | 9.2×10^{-8} | 75,884 | 0.37 |
| Component 2: UKBB | 0.51 | 0.029 | 0.005 | 7.6×10^{-8} | 67,786 | N/A |
| European ancestry meta-analysis | 0.51 | 0.028 | 0.004 | 3.0×10^{-14} | 143,670 | 0.81 |
| Component 3: non-European ancestry | 0.77 | 0.025 | 0.017 | 0.13 | 10,103 | 0.12 |
| Trans-ancestry meta-analysis | 0.52 | 0.028 | 0.004 | 9.5×10^{-15} | 153,773 | 0.80 |

Locus: *NT5C2*. Lead SNP: rs74233809. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.08 | 0.032 | 0.009 | 0.00039 | 75,881 | 0.88 |
| Component 2: UKBB | 0.08 | 0.042 | 0.010 | 3.0×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.08 | 0.037 | 0.007 | 5.2×10^{-8} | 143,667 | 0.49 |
| Component 3: non-European ancestry | 0.12 | 0.061 | 0.022 | 0.0057 | 10,104 | 0.39 |
| Trans-ancestry meta-analysis | 0.08 | 0.039 | 0.006 | 1.8×10^{-9} | 153,771 | 0.14 |

Locus: *ADRB1*. Lead SNP: rs7076938. Effect/other alleles: T/C.

| Analysis | EAf | B | SE | P-value | N | Q P-value |
|------------------------------------|------|-------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.73 | 0.046 | 0.006 | 1.3×10^{-15} | 75,885 | 0.79 |
| Component 2: UKBB | 0.73 | 0.025 | 0.006 | 4.5×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.73 | 0.036 | 0.004 | 4.7×10^{-18} | 143,671 | 0.011 |
| Component 3: non-European ancestry | 0.64 | 0.018 | 0.015 | 0.22 | 10,103 | 0.56 |
| Trans-ancestry meta-analysis | 0.73 | 0.035 | 0.004 | 4.7×10^{-18} | 153,774 | 0.19 |

Locus: *PLEKHA1*. Lead SNP: rs2421016. Effect/other alleles: T/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.50 | 0.022 | 0.005 | 1.0×10^{-5} | 75,873 | 0.25 |
| Component 2: UKBB | 0.47 | 0.019 | 0.005 | 0.00047 | 67,786 | N/A |
| European ancestry meta-analysis | 0.49 | 0.021 | 0.004 | 1.8×10^{-8} | 143,659 | 0.62 |
| Component 3: non-European ancestry | 0.40 | 0.021 | 0.014 | 0.15 | 10,100 | 0.28 |
| Trans-ancestry meta-analysis | 0.48 | 0.021 | 0.004 | 6.1×10^{-9} | 153,759 | 0.83 |

Locus: *INS-IGF2*. Lead SNP: rs72851023. Effect/other alleles: T/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.07 | 0.055 | 0.011 | 6.7×10^{-7} | 67,990 | 0.014 |
| Component 2: UKBB | 0.08 | 0.041 | 0.010 | 8.0×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.07 | 0.048 | 0.008 | 2.9×10^{-10} | 135,776 | 0.35 |
| Component 3: non-European ancestry | 0.02 | -0.048 | 0.064 | 0.45 | 7,815 | 1.00 |
| Trans-ancestry meta-analysis | 0.07 | 0.046 | 0.007 | 6.8×10^{-10} | 143,591 | 0.64 |

Locus: *MTNR1B*. Lead SNP: rs10830963. Effect/other alleles: G/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.28 | 0.029 | 0.006 | 5.1×10^{-7} | 75,877 | 0.20 |
| Component 2: UKBB | 0.28 | 0.017 | 0.006 | 0.0059 | 67,786 | N/A |
| European ancestry meta-analysis | 0.28 | 0.023 | 0.004 | 2.9×10^{-8} | 143,663 | 0.12 |
| Component 3: non-European ancestry | 0.20 | -0.013 | 0.021 | 0.54 | 10,102 | 0.76 |
| Trans-ancestry meta-analysis | 0.27 | 0.022 | 0.004 | 1.0×10^{-7} | 153,765 | 0.23 |

Locus: *APOLD1*. Lead SNP: rs11055034. Effect/other alleles: C/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.74 | 0.028 | 0.006 | 2.1×10^{-6} | 75,867 | 0.91 |
| Component 2: UKBB | 0.72 | 0.016 | 0.006 | 0.0099 | 67,786 | N/A |
| European ancestry meta-analysis | 0.73 | 0.022 | 0.004 | 1.8×10^{-7} | 143,653 | 0.13 |
| Component 3: non-European ancestry | 0.81 | 0.046 | 0.020 | 0.019 | 9,942 | 0.67 |
| Trans-ancestry meta-analysis | 0.73 | 0.023 | 0.004 | 2.3×10^{-8} | 153,595 | 0.32 |

Locus: *ABCC9*. Lead SNP: rs139975827. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.61 | 0.027 | 0.007 | 4.9×10^{-5} | 55,417 | 0.72 |
| Component 2: UKBB | 0.63 | 0.023 | 0.006 | 5.8×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.62 | 0.025 | 0.004 | 1.1×10^{-8} | 123,203 | 0.72 |
| Component 3: non-European ancestry | 0.70 | -0.018 | 0.018 | 0.32 | 8,655 | 0.68 |
| Trans-ancestry meta-analysis | 0.63 | 0.022 | 0.004 | 1.0×10^{-7} | 131,858 | 0.22 |

Locus: *ITPR2*. Lead SNP: rs12823128. Effect/other alleles: T/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.54 | 0.017 | 0.005 | 0.00086 | 71,645 | 0.59 |
| Component 2: UKBB | 0.53 | 0.025 | 0.005 | 3.8×10^{-6} | 67,786 | N/A |
| European ancestry meta-analysis | 0.54 | 0.021 | 0.004 | 1.9×10^{-8} | 139,431 | 0.32 |
| Component 3: non-European ancestry | 0.86 | 0.000 | 0.021 | 1.0 | 10,103 | 0.054 |
| Trans-ancestry meta-analysis | 0.56 | 0.020 | 0.004 | 3.2×10^{-8} | 149,534 | 0.65 |

Locus: *HMGA2*. Lead SNP: rs1351394. Effect/other alleles: T/C.

| Analysis | EAf | B | SE | P-value | N | Q P-value |
|------------------------------------|------|-------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.50 | 0.048 | 0.005 | 7.6×10^{-21} | 75,885 | 0.40 |
| Component 2: UKBB | 0.49 | 0.039 | 0.005 | 3.1×10^{-13} | 67,786 | N/A |
| European ancestry meta-analysis | 0.49 | 0.044 | 0.004 | 1.9×10^{-32} | 143,671 | 0.24 |
| Component 3: non-European ancestry | 0.34 | 0.033 | 0.015 | 0.028 | 10,102 | 0.75 |
| Trans-ancestry meta-analysis | 0.48 | 0.043 | 0.004 | 2.0×10^{-33} | 153,773 | 0.72 |

Locus: *IGF1*. Lead SNP: rs7964361. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.08 | 0.038 | 0.009 | 5.6×10^{-5} | 71,642 | 0.63 |
| Component 2: UKBB | 0.09 | 0.040 | 0.010 | 2.3×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.09 | 0.039 | 0.007 | 4.7×10^{-9} | 139,428 | 0.86 |
| Component 3: non-European ancestry | 0.03 | -0.010 | 0.040 | 0.81 | 9,264 | 0.21 |
| Trans-ancestry meta-analysis | 0.08 | 0.038 | 0.007 | 9.7×10^{-9} | 148,692 | 0.20 |

Locus: *LINC00332*. Lead SNP: rs2324499. Effect/other alleles: G/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.68 | 0.026 | 0.006 | 7.9×10^{-6} | 71,639 | 0.25 |
| Component 2: UKBB | 0.67 | 0.018 | 0.006 | 0.0019 | 67,786 | N/A |
| European ancestry meta-analysis | 0.68 | 0.022 | 0.004 | 7.3×10^{-8} | 139,425 | 0.34 |
| Component 3: non-European ancestry | 0.60 | 0.034 | 0.016 | 0.027 | 8,655 | 0.41 |
| Trans-ancestry meta-analysis | 0.67 | 0.023 | 0.004 | 8.3×10^{-9} | 148,080 | 0.64 |

Locus: *RB1*. Lead SNP: rs2854355. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.28 | 0.024 | 0.006 | 0.00012 | 69,775 | 0.036 |
| Component 2: UKBB | 0.26 | 0.023 | 0.006 | 0.00024 | 67,786 | N/A |
| European ancestry meta-analysis | 0.27 | 0.023 | 0.004 | 9.8×10^{-8} | 137,561 | 0.91 |
| Component 3: non-European ancestry | 0.19 | 0.037 | 0.020 | 0.066 | 8,654 | 0.86 |
| Trans-ancestry meta-analysis | 0.26 | 0.024 | 0.004 | 2.2×10^{-8} | 146,215 | 0.63 |

Locus: *RNF219-AS1*. Lead SNP: rs1819436. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.86 | 0.041 | 0.008 | 2.4×10^{-7} | 71,193 | 0.19 |
| Component 2: UKBB | 0.88 | 0.024 | 0.008 | 0.0030 | 67,786 | N/A |
| European ancestry meta-analysis | 0.87 | 0.033 | 0.006 | 6.3×10^{-9} | 138,979 | 0.14 |
| Component 3: non-European ancestry | 0.84 | 0.033 | 0.021 | 0.12 | 8,654 | 0.96 |
| Trans-ancestry meta-analysis | 0.87 | 0.033 | 0.005 | 1.8×10^{-9} | 147,633 | 0.80 |

Locus: *FES*. Lead SNP: rs12906125. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.69 | 0.020 | 0.006 | 0.00036 | 73,495 | 0.92 |
| Component 2: UKBB | 0.67 | 0.025 | 0.006 | 1.1×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.68 | 0.023 | 0.004 | 1.7×10^{-8} | 141,281 | 0.55 |
| Component 3: non-European ancestry | 0.75 | 0.018 | 0.017 | 0.30 | 10,103 | 0.87 |
| Trans-ancestry meta-analysis | 0.69 | 0.023 | 0.004 | 1.0×10^{-8} | 151,384 | 0.75 |

Locus: *IGF1R*. Lead SNP: rs7402982. Effect/other alleles: A/G.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.43 | 0.022 | 0.005 | 4.9×10^{-5} | 71,637 | 0.24 |
| Component 2: UKBB | 0.41 | 0.024 | 0.006 | 1.3×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.43 | 0.023 | 0.004 | 2.3×10^{-9} | 139,423 | 0.83 |
| Component 3: non-European ancestry | 0.35 | 0.020 | 0.017 | 0.23 | 10,103 | 0.61 |
| Trans-ancestry meta-analysis | 0.42 | 0.023 | 0.004 | 1.1×10^{-9} | 149,526 | 0.61 |

Locus: *GPR139*. Lead SNP: rs1011939. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.31 | 0.022 | 0.006 | 0.00013 | 75,818 | 0.93 |
| Component 2: UKBB | 0.28 | 0.022 | 0.006 | 0.00030 | 67,786 | N/A |
| European ancestry meta-analysis | 0.30 | 0.022 | 0.004 | 1.3×10^{-7} | 143,604 | 0.97 |
| Component 3: non-European ancestry | 0.57 | 0.047 | 0.015 | 0.0013 | 10,098 | 0.52 |
| Trans-ancestry meta-analysis | 0.31 | 0.024 | 0.004 | 2.7×10^{-9} | 153,702 | 0.66 |

Locus: *CLDN7*. Lead SNP: rs113086489. Effect/other alleles: T/C.

| Analysis | EAF | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.57 | 0.033 | 0.005 | 7.1×10^{-10} | 71,640 | 0.10 |
| Component 2: UKBB | 0.55 | 0.028 | 0.005 | 2.6×10^{-7} | 67,786 | N/A |
| European ancestry meta-analysis | 0.56 | 0.031 | 0.004 | 9.1×10^{-16} | 139,426 | 0.51 |
| Component 3: non-European ancestry | 0.40 | 0.012 | 0.015 | 0.40 | 10,104 | 0.29 |
| Trans-ancestry meta-analysis | 0.55 | 0.030 | 0.004 | 1.3×10^{-15} | 149,530 | 0.77 |

Locus: *SUZ12P1-CRLF3*. Lead SNP: rs144843919. Effect/other alleles: G/A.

| Analysis | EAF | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.97 | 0.036 | 0.018 | 0.0048 | 53,571 | 0.13 |
| Component 2: UKBB | 0.96 | 0.087 | 0.015 | 1.0×10^{-8} | 67,786 | N/A |
| European ancestry meta-analysis | 0.96 | 0.066 | 0.012 | 1.4×10^{-8} | 121,357 | 0.033 |
| Component 3: non-European ancestry | 0.96 | 0.112 | 0.049 | 0.022 | 6,635 | 0.056 |
| Trans-ancestry meta-analysis | 0.96 | 0.068 | 0.011 | 1.5×10^{-9} | 127,992 | 0.068 |

Locus: *SP6-SP2*. Lead SNP: rs12942207. Effect/other alleles: C/T.

| Analysis | EAF | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.31 | 0.025 | 0.006 | 2.2×10^{-5} | 71,635 | 0.85 |
| Component 2: UKBB | 0.30 | 0.020 | 0.006 | 0.00063 | 67,786 | N/A |
| European ancestry meta-analysis | 0.30 | 0.022 | 0.004 | 5.1×10^{-8} | 139,421 | 0.59 |
| Component 3: non-European ancestry | 0.26 | 0.050 | 0.018 | 0.0053 | 8,654 | 0.98 |
| Trans-ancestry meta-analysis | 0.30 | 0.024 | 0.004 | 3.0×10^{-9} | 148,075 | 0.75 |

Locus: *ACTL9*. Lead SNP: rs61154119. Effect/other alleles: T/G.

| Analysis | EAF | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.84 | 0.030 | 0.007 | 6.4×10^{-5} | 63,412 | 0.00014 |
| Component 2: UKBB | 0.85 | 0.026 | 0.007 | 0.00050 | 67,786 | N/A |
| European ancestry meta-analysis | 0.84 | 0.028 | 0.005 | 1.1×10^{-7} | 131,198 | 0.70 |
| Component 3: non-European ancestry | 0.79 | 0.033 | 0.018 | 0.077 | 8,655 | 0.64 |
| Trans-ancestry meta-analysis | 0.84 | 0.028 | 0.005 | 2.3×10^{-8} | 139,853 | 0.98 |

Locus: *PEPD*. Lead SNP: rs10402712. Effect/other alleles: A/G.

| Analysis | EAF | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.25 | 0.021 | 0.006 | 0.00033 | 71,642 | 0.23 |
| Component 2: UKBB | 0.26 | 0.022 | 0.006 | 0.00042 | 67,786 | N/A |
| European ancestry meta-analysis | 0.26 | 0.022 | 0.004 | 4.4×10^{-7} | 139,428 | 0.98 |
| Component 3: non-European ancestry | 0.40 | 0.038 | 0.014 | 0.0086 | 10,102 | 0.89 |
| Trans-ancestry meta-analysis | 0.27 | 0.023 | 0.004 | 2.3×10^{-8} | 149,530 | 0.72 |

Locus: *JAG1*. Lead SNP: rs6040076. Effect/other alleles: C/G.

| Analysis | EAF | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.48 | 0.023 | 0.005 | 3.1×10^{-5} | 71,638 | 0.82 |
| Component 2: UKBB | 0.51 | 0.023 | 0.005 | 1.8×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.49 | 0.023 | 0.004 | 2.0×10^{-9} | 139,424 | 0.96 |
| Component 3: non-European ancestry | 0.69 | -0.006 | 0.018 | 0.73 | 8,655 | 0.026 |
| Trans-ancestry meta-analysis | 0.51 | 0.022 | 0.004 | 7.2×10^{-9} | 148,079 | 0.24 |

Locus: *C20orf203*. Lead SNP: rs28530618. Effect/other alleles: A/G.

| Analysis | EAF | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.50 | 0.032 | 0.005 | 5.0×10^{-9} | 70,376 | 0.45 |
| Component 2: UKBB | 0.47 | 0.020 | 0.005 | 0.00015 | 67,786 | N/A |
| European ancestry meta-analysis | 0.49 | 0.026 | 0.004 | 7.7×10^{-12} | 138,162 | 0.14 |
| Component 3: non-European ancestry | 0.63 | -0.011 | 0.016 | 0.50 | 10,102 | 0.25 |
| Trans-ancestry meta-analysis | 0.50 | 0.024 | 0.004 | 8.4×10^{-11} | 148,264 | 0.18 |

Locus: *MAFB*. Lead SNP: rs6016377. Effect/other alleles: T/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.41 | 0.028 | 0.005 | 3.8×10^{-7} | 71,639 | 0.10 |
| Component 2: UKBB | 0.45 | 0.020 | 0.006 | 0.00043 | 67,786 | N/A |
| European ancestry meta-analysis | 0.43 | 0.024 | 0.004 | 9.5×10^{-10} | 139,425 | 0.29 |
| Component 3: non-European ancestry | 0.74 | 0.025 | 0.018 | 0.17 | 10,103 | 0.26 |
| Trans-ancestry meta-analysis | 0.45 | 0.024 | 0.004 | 3.7×10^{-10} | 149,528 | 0.27 |

Locus: *NRIP1*. Lead SNP: rs2229742. Effect/other alleles: G/C.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.85 | 0.037 | 0.008 | 6.6×10^{-6} | 75,886 | 0.66 |
| Component 2: UKBB | 0.90 | 0.034 | 0.009 | 9.1×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.87 | 0.036 | 0.006 | 2.2×10^{-9} | 143,672 | 0.82 |
| Component 3: non-European ancestry | 0.82 | -0.085 | 0.044 | 0.052 | 9,264 | 0.72 |
| Trans-ancestry meta-analysis | 0.87 | 0.034 | 0.006 | 1.5×10^{-8} | 152,936 | 0.092 |

Locus: *KREMEN1*. Lead SNP: rs134594. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|---------|-----------|
| Component 1: European ancestry | 0.35 | 0.022 | 0.006 | 8.9×10^{-5} | 69,554 | 0.36 |
| Component 2: UKBB | 0.35 | 0.023 | 0.006 | 3.2×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.35 | 0.023 | 0.004 | 1.0×10^{-8} | 137,340 | 0.84 |
| Component 3: non-European ancestry | 0.33 | 0.002 | 0.016 | 0.90 | 8,655 | 0.59 |
| Trans-ancestry meta-analysis | 0.35 | 0.022 | 0.004 | 2.2×10^{-8} | 145,995 | 0.25 |

Locus: *SREBF2*. Lead SNP: rs62240962. Effect/other alleles: C/T.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|-----------------------|---------|-----------|
| Component 1: European ancestry | 0.91 | 0.054 | 0.010 | 2.6×10^{-8} | 68,286 | 0.69 |
| Component 2: UKBB | 0.92 | 0.040 | 0.010 | 6.0×10^{-5} | 67,786 | N/A |
| European ancestry meta-analysis | 0.91 | 0.047 | 0.007 | 9.7×10^{-12} | 136,072 | 0.30 |
| Component 3: non-European ancestry | 0.97 | 0.07 | 0.048 | 0.15 | 7,815 | 0.028 |
| Trans-ancestry meta-analysis | 0.92 | 0.047 | 0.007 | 3.7×10^{-12} | 143,887 | 0.69 |

Locus: *PLAC1*. Lead SNP: rs11096402. Effect/other alleles: G/A.

| Analysis | EAf | β | SE | P-value | N | Q P-value |
|------------------------------------|------|---------|-------|----------------------|--------|-----------|
| Component 1: European ancestry | 0.28 | 0.027 | 0.007 | 2.0×10^{-4} | 37,417 | 0.73 |
| Component 2: UKBB | 0.24 | 0.028 | 0.006 | 1.8×10^{-6} | 57,620 | N/A |
| European ancestry meta-analysis | 0.25 | 0.028 | 0.005 | 1.3×10^{-9} | 95,037 | 0.31 |
| Component 3: non-European ancestry | N/A | N/A | N/A | N/A | N/A | N/A |
| Trans-ancestry meta-analysis | N/A | N/A | N/A | N/A | N/A | N/A |

PLAC1 result (on X chromosome) was not available in non-European ancestry studies.

EAf, effect allele frequency; SE, standard error.

Supplementary Table 4. Summary statistics for lead SNPs attaining genome-wide significant evidence ($P < 5 \times 10^{-8}$) of association with BW in European ancestry meta-analysis of up to 143,677 individuals and/or trans-ancestry meta-analysis of up to 153,781 individuals.

| Locus | SNP | Chr | Position (b37, bp) | Alleles | | EAF | European ancestry | | Trans-ancestry | | Status ^a | EUR r^2 with lead TA | Present in HapMap? | Best proxy in HapMap | |
|--------------------|-------------|-----|-----------------------|---------|-------|------|-------------------|-----------------------|------------------|-----------------------|---------------------|---------------------------|-----------------------|----------------------|-----------|
| | | | | Effect | Other | | β (SE) | P-value | β (SE) | P-value | | | | SNP | EUR r^2 |
| <i>WNT4-ZBTB40</i> | rs2473248 | 1 | 22,536,643 | C | T | 0.87 | 0.033 (0.006) | 1.1×10^{-8} | 0.033 (0.005) | 1.1×10^{-9} | TA/EUR | | No | rs2744726 | 0.99 |
| <i>ZBTB7B</i> | rs3753639 | 1 | 154,986,091 | C | T | 0.24 | 0.031 (0.004) | 7.3×10^{-12} | 0.031 (0.004) | 1.3×10^{-12} | TA/EUR | | No | rs905938 | 0.77 |
| <i>FCGR2B</i> | rs72480273 | 1 | 161,644,871 | C | A | 0.17 | 0.031 (0.005) | 8.0×10^{-10} | 0.030 (0.005) | 1.5×10^{-9} | TA/EUR | | No | rs17413015 | 0.97 |
| <i>DTL</i> | rs61830764 | 1 | 212,289,976 | A | G | 0.38 | 0.022 (0.004) | 5.6×10^{-8} | 0.022 (0.004) | 4.5×10^{-8} | TA | | No | rs1387815 | 0.51 |
| <i>ATAD2B</i> | rs7575873 | 2 | 23,962,647 | A | G | 0.88 | 0.038 (0.006) | 1.3×10^{-11} | 0.036 (0.006) | 6.2×10^{-11} | TA | | No | rs718139 | 0.98 |
| | rs181211713 | 2 | 23,965,445 | G | A | 0.87 | 0.038 (0.006) | 1.2×10^{-11} | 0.036 (0.005) | 7.3×10^{-11} | EUR | 0.94 | No | | |
| <i>EPAS1</i> | rs1374204 | 2 | 46,484,205 | T | C | 0.70 | 0.047 (0.004) | 6.2×10^{-29} | 0.046 (0.004) | 1.5×10^{-29} | TA | | Yes | | |
| | rs17034876 | 2 | 46,484,310 | T | C | 0.70 | 0.047 (0.004) | 2.6×10^{-29} | 0.046 (0.004) | 1.9×10^{-29} | EUR | 0.99 | Yes | | |
| <i>PTH1R</i> | rs2242116 | 3 | 46,941,116 | A | G | 0.38 | 0.022 (0.004) | 1.4×10^{-8} | 0.021 (0.004) | 1.2×10^{-8} | TA | | Yes | | |
| | rs2168443 | 3 | 46,947,087 | T | A | 0.38 | 0.023 (0.004) | 3.5×10^{-9} | 0.021 (0.004) | 1.3×10^{-8} | EUR | 0.99 | Yes | | |
| <i>ADCY5</i> | rs11719201 | 3 | 123,068,744 | T | C | 0.23 | 0.046 (0.004) | 2.4×10^{-26} | 0.046 (0.004) | 6.4×10^{-27} | TA/EUR | | No | rs7613951 | 0.97 |
| | rs9883204 | 3 | 123,096,820 | T | C | 0.25 | 0.041 (0.004) | 8.4×10^{-22} | 0.040 (0.004) | 6.1×10^{-22} | Previous | 0.85 | Yes | | |
| <i>CPA3</i> | rs10935733 | 3 | 148,622,968 | T | C | 0.41 | 0.022 (0.004) | 9.2×10^{-9} | 0.023 (0.004) | 6.2×10^{-10} | TA/EUR | | No | rs12496367 | 0.81 |
| <i>CCNL1-LEKR1</i> | rs13322435 | 3 | 156,795,468 | A | G | 0.60 | 0.053 (0.004) | 3.7×10^{-41} | 0.052 (0.004) | 1.3×10^{-42} | TA | | Yes | | |
| | rs900399 | 3 | 156,798,732 | A | G | 0.61 | 0.052 (0.004) | 2.2×10^{-41} | 0.051 (0.004) | 2.9×10^{-41} | EUR | 0.92 | Yes | | |
| | rs900400 | 3 | 156,798,775 | T | C | 0.61 | 0.053 (0.004) | 1.1×10^{-40} | 0.051 (0.004) | 8.4×10^{-41} | Previous | 0.92 | Yes | | |
| <i>LCORL</i> | rs925098 | 4 | 17,919,811 | G | A | 0.27 | 0.034 (0.004) | 5.4×10^{-16} | 0.032 (0.004) | 1.3×10^{-15} | TA | | Yes | | |
| | rs2724475 | 4 | 17,946,432 | T | C | 0.27 | 0.034 (0.004) | 5.3×10^{-16} | 0.032 (0.004) | 1.3×10^{-15} | EUR | 0.97 | Yes | | |
| | rs724577 | 4 | 17,993,410 | A | C | 0.27 | 0.033 (0.004) | 1.7×10^{-15} | 0.031 (0.004) | 1.5×10^{-14} | Previous | 0.98 | Yes | | |

| Locus | SNP | Chr | Position (b37, bp) | Alleles | | EAF | European ancestry | | Trans-ancestry | | Status ^a | EUR r^2 with lead TA | Present in HapMap? | Best proxy in HapMap | |
|-----------|-------------|-----|-----------------------|---------|-------|------|-------------------|-----------------------|------------------|-----------------------|---------------------|---------------------------|-----------------------|----------------------|-----------|
| | | | | Effect | Other | | β (SE) | P-value | β (SE) | P-value | | | | SNP | EUR r^2 |
| HHIP | rs6537307 | 4 | 145,601,863 | G | A | 0.50 | 0.025 (0.004) | 9.5×10^{-12} | 0.026 (0.004) | 1.3×10^{-12} | TA | | Yes | | |
| | rs2131354 | 4 | 145,599,908 | A | G | 0.53 | 0.026 (0.004) | 4.1×10^{-12} | 0.026 (0.004) | 2.3×10^{-12} | EUR | 0.88 | No | | |
| 5q11.2 | rs854037 | 5 | 57,091,783 | A | G | 0.81 | 0.027 (0.005) | 2.2×10^{-8} | 0.025 (0.005) | 3.5×10^{-8} | TA/EUR | | Yes | | |
| | rs4432842 | 5 | 57,172,078 | T | C | 0.70 | 0.021 (0.004) | 3.4×10^{-7} | 0.019 (0.004) | 7.0×10^{-7} | Previous | 0.47 | Yes | | |
| EBF1 | rs7729301 | 5 | 157,886,953 | A | G | 0.73 | 0.024 (0.004) | 1.6×10^{-8} | 0.025 (0.004) | 1.3×10^{-9} | TA | | No | rs2946164 | 0.97 |
| | rs2946179 | 5 | 157,886,627 | C | T | 0.73 | 0.024 (0.004) | 1.3×10^{-8} | 0.024 (0.004) | 1.5×10^{-9} | EUR | 0.98 | No | | |
| CDKAL1 | rs35261542 | 6 | 20,675,792 | C | A | 0.73 | 0.044 (0.004) | 4.4×10^{-27} | 0.044 (0.004) | 9.7×10^{-29} | TA/EUR | | No | rs7756992 | 0.97 |
| | rs6931514 | 6 | 20,703,952 | A | G | 0.73 | 0.043 (0.004) | 6.2×10^{-26} | 0.043 (0.004) | 1.0×10^{-26} | Previous | 0.94 | Yes | | |
| HIST1H2BE | rs9379832 | 6 | 26,186,200 | A | G | 0.70 | 0.023 (0.004) | 6.6×10^{-8} | 0.024 (0.004) | 1.2×10^{-8} | TA | | No | rs806794 | 0.62 |
| HMGA1 | rs7742369 | 6 | 34,165,721 | G | A | 0.18 | 0.028 (0.005) | 1.0×10^{-8} | 0.027 (0.005) | 1.1×10^{-8} | TA | | Yes | | |
| | rs1187118 | 6 | 34,169,020 | A | T | 0.17 | 0.030 (0.005) | 3.6×10^{-9} | 0.027 (0.005) | 1.4×10^{-8} | EUR | 0.95 | No | | |
| L3MBTL3 | rs1415701 | 6 | 130,345,835 | G | A | 0.74 | 0.025 (0.004) | 2.6×10^{-9} | 0.027 (0.004) | 4.0×10^{-11} | TA/EUR | | Yes | | |
| ESR1 | rs1101081 | 6 | 152,032,917 | C | T | 0.72 | 0.038 (0.004) | 1.6×10^{-19} | 0.037 (0.004) | 6.1×10^{-20} | TA | | No | rs851977 | 0.99 |
| | rs10872678 | 6 | 152,039,964 | T | C | 0.72 | 0.038 (0.004) | 6.9×10^{-20} | 0.036 (0.004) | 1.3×10^{-19} | EUR | 0.98 | Yes | | |
| GNA12 | rs798489 | 7 | 2,801,803 | C | T | 0.73 | 0.023 (0.004) | 2.0×10^{-8} | 0.024 (0.004) | 5.0×10^{-9} | TA | | Yes | | |
| | rs798498 | 7 | 2,795,882 | T | G | 0.69 | 0.023 (0.004) | 1.3×10^{-8} | 0.022 (0.004) | 3.8×10^{-8} | EUR | 0.83 | Yes | | |
| IGF2BP3 | rs11765649 | 7 | 23,479,013 | T | C | 0.75 | 0.027 (0.004) | 5.8×10^{-10} | 0.026 (0.004) | 1.0×10^{-9} | TA/EUR | | No | rs12540730 | 0.86 |
| TBX20 | rs6959887 | 7 | 35,295,365 | A | G | 0.61 | 0.023 (0.004) | 1.5×10^{-9} | 0.021 (0.004) | 1.0×10^{-8} | TA/EUR | | Yes | | |
| YKT6-GCK | rs138715366 | 7 | 44,246,271 | C | T | 0.99 | 0.241 (0.023) | 7.2×10^{-26} | 0.244 (0.023) | 1.4×10^{-26} | TA/EUR | | No | rs2908277 | 0.096 |
| MLXIPL | rs62466330 | 7 | 73,056,805 | C | T | 0.07 | 0.049 (0.008) | 1.2×10^{-10} | 0.051 (0.007) | 5.9×10^{-12} | TA | | No | rs17401675 | 1 |
| | rs111778406 | 7 | 72,957,570 | G | A | 0.07 | 0.049 (0.008) | 5.8×10^{-11} | 0.050 (0.007) | 1.3×10^{-11} | EUR | 0.78 | No | | |

| Locus | SNP | Chr | Position | Alleles | | EAF | European ancestry | | Trans-ancestry | | Status ^a | EUR r^2 with lead TA | Present in HapMap? | Best proxy in HapMap | |
|--------------------|------------|-----|-------------|---------|-------|------|-------------------|-----------------------|----------------|-----------------------|---------------------|------------------------|--------------------|----------------------|-----------|
| | | | (b37, bp) | Effect | Other | | β (SE) | P-value | β (SE) | P-value | | | | SNP | EUR r^2 |
| <i>ANK1-NKX6.3</i> | rs13266210 | 8 | 41,533,514 | A | G | 0.79 | 0.031 (0.005) | 1.3×10^{-11} | 0.030 (0.004) | 1.6×10^{-11} | TA/EUR | | Yes | | |
| <i>TRIB1</i> | rs6989280 | 8 | 126,508,746 | G | A | 0.73 | 0.022 (0.004) | 2.2×10^{-7} | 0.022 (0.004) | 5.0×10^{-8} | TA | | Yes | | |
| <i>SLC45A4</i> | rs12543725 | 8 | 142,247,979 | G | A | 0.59 | 0.023 (0.004) | 1.2×10^{-9} | 0.022 (0.004) | 1.9×10^{-9} | TA/EUR | | Yes | | |
| <i>PTCH1</i> | rs28510415 | 9 | 98,245,026 | G | A | 0.09 | 0.056 (0.007) | 1.5×10^{-17} | 0.053 (0.006) | 4.0×10^{-16} | TA | | No | rs3824488 | 1 |
| | rs3780573 | 9 | 98,239,503 | A | G | 0.10 | 0.055 (0.006) | 7.0×10^{-18} | 0.050 (0.006) | 1.1×10^{-15} | EUR | 1.00 | No | | |
| <i>LPAR1</i> | rs2150052 | 9 | 113,945,067 | T | A | 0.50 | 0.021 (0.004) | 2.2×10^{-8} | 0.020 (0.004) | 2.8×10^{-8} | TA | | Yes | | |
| | rs1411424 | 9 | 113,892,963 | A | G | 0.52 | 0.021 (0.004) | 2.2×10^{-8} | 0.020 (0.004) | 4.1×10^{-8} | EUR | 0.86 | Yes | | |
| <i>PHF19</i> | rs7847628 | 9 | 123,631,225 | G | A | 0.68 | 0.023 (0.004) | 1.0×10^{-8} | 0.023 (0.004) | 5.4×10^{-9} | TA | | Yes | rs1056567 | 0.87 |
| | rs4836833 | 9 | 123,632,829 | C | G | 0.67 | 0.023 (0.004) | 8.6×10^{-9} | 0.022 (0.004) | 1.5×10^{-8} | EUR | 0.88 | Yes | | |
| <i>STRBP</i> | rs700059 | 9 | 125,824,055 | G | A | 0.14 | 0.033 (0.005) | 4.7×10^{-10} | 0.036 (0.005) | 1.2×10^{-12} | TA | | Yes | | |
| | rs10818797 | 9 | 126,020,405 | C | T | 0.14 | 0.035 (0.005) | 1.2×10^{-10} | 0.035 (0.005) | 1.5×10^{-11} | EUR | 0.91 | Yes | | |
| <i>HHEX-IDE</i> | rs61862780 | 10 | 94,468,643 | T | C | 0.51 | 0.028 (0.004) | 3.0×10^{-14} | 0.028 (0.004) | 9.5×10^{-15} | TA | | No | rs2497306 | 0.97 |
| | rs2497304 | 10 | 94,492,716 | C | T | 0.52 | 0.028 (0.004) | 2.6×10^{-14} | 0.028 (0.004) | 1.4×10^{-14} | EUR | 0.90 | Yes | | |
| <i>NT5C2</i> | rs74233809 | 10 | 104,913,940 | C | T | 0.08 | 0.037 (0.007) | 5.2×10^{-8} | 0.039 (0.006) | 1.8×10^{-9} | TA | | No | rs11191582 | 1 |
| | rs79237883 | 10 | 104,940,946 | C | T | 0.08 | 0.037 (0.007) | 3.6×10^{-8} | 0.038 (0.006) | 5.6×10^{-9} | EUR | 1.00 | No | | |
| <i>ADRB1</i> | rs7076938 | 10 | 115,789,375 | T | C | 0.73 | 0.036 (0.004) | 4.7×10^{-18} | 0.035 (0.004) | 4.7×10^{-18} | TA | | Yes | | |
| | rs740746 | 10 | 115,792,787 | A | G | 0.73 | 0.036 (0.004) | 3.8×10^{-18} | 0.035 (0.004) | 6.4×10^{-18} | EUR | 0.99 | Yes | | |
| | rs1801253 | 10 | 115,805,056 | C | G | 0.71 | 0.032 (0.004) | 3.0×10^{-14} | 0.031 (0.004) | 2.6×10^{-14} | Previous | 0.99 | Yes | | |
| <i>PLEKHA1</i> | rs2421016 | 10 | 124,167,512 | T | C | 0.49 | 0.021 (0.004) | 1.8×10^{-8} | 0.021 (0.004) | 6.1×10^{-9} | TA/EUR | | Yes | | |
| <i>INS-IGF2</i> | rs72851023 | 11 | 2,130,620 | T | C | 0.07 | 0.048 (0.008) | 2.9×10^{-10} | 0.046 (0.007) | 6.8×10^{-10} | TA/EUR | | No | rs868332 | 0.49 |
| <i>MTNR1B</i> | rs10830963 | 11 | 92,708,710 | G | C | 0.28 | 0.023 (0.004) | 2.9×10^{-8} | 0.022 (0.004) | 1.0×10^{-7} | EUR | | Yes | | |

| Locus | SNP | Chr | Position | Alleles | | EAF | European ancestry | | Trans-ancestry | | Status ^a | EUR r^2 with lead TA | Present in HapMap? | Best proxy in HapMap | |
|---------------|-------------|-----|-------------|---------|-------|------|-------------------|-----------------------|----------------|-----------------------|---------------------|------------------------|--------------------|----------------------|-----------|
| | | | (b37, bp) | Effect | Other | | β (SE) | P-value | β (SE) | P-value | | | | SNP | EUR r^2 |
| APOLD1 | rs11055034 | 12 | 12,890,626 | C | A | 0.73 | 0.022 (0.004) | 1.8×10^{-7} | 0.023 (0.004) | 2.3×10^{-8} | TA | | Yes | | |
| ABCC9 | rs139975827 | 12 | 22,068,161 | G | A | 0.62 | 0.025 (0.004) | 1.1×10^{-8} | 0.022 (0.004) | 1.0×10^{-7} | EUR | | No | rs4148656 | 0.67 |
| ITPR2 | rs12823128 | 12 | 26,872,730 | T | C | 0.54 | 0.021 (0.004) | 1.9×10^{-8} | 0.020 (0.004) | 3.2×10^{-8} | TA | | Yes | | |
| | rs2306547 | 12 | 26,877,885 | C | T | 0.54 | 0.021 (0.004) | 1.8×10^{-8} | 0.020 (0.004) | 3.2×10^{-8} | EUR | 0.96 | Yes | | |
| HMGA2 | rs1351394 | 12 | 66,351,826 | T | C | 0.49 | 0.044 (0.004) | 1.9×10^{-32} | 0.043 (0.004) | 2.0×10^{-33} | TA/EUR | | Yes | | |
| | rs1042725 | 12 | 66,358,347 | C | T | 0.51 | 0.043 (0.004) | 3.1×10^{-32} | 0.042 (0.004) | 7.1×10^{-32} | Previous | 0.93 | Yes | | |
| IGF1 | rs7964361 | 12 | 102,994,878 | A | G | 0.09 | 0.039 (0.007) | 4.7×10^{-9} | 0.038 (0.007) | 9.7×10^{-9} | TA/EUR | | Yes | | |
| LINCO0332 | rs2324499 | 13 | 40,662,001 | G | C | 0.68 | 0.022 (0.004) | 7.3×10^{-8} | 0.023 (0.004) | 8.3×10^{-9} | TA | | Yes | | |
| | rs7998537 | 13 | 40,662,742 | G | A | 0.68 | 0.022 (0.004) | 3.9×10^{-8} | 0.022 (0.004) | 1.8×10^{-8} | EUR | 1.00 | Yes | | |
| RB1 | rs2854355 | 13 | 48,882,363 | G | A | 0.27 | 0.023 (0.004) | 9.8×10^{-8} | 0.024 (0.004) | 2.2×10^{-8} | TA | | No | rs9568028 | 0.88 |
| | rs34217484 | 13 | 48,854,550 | A | T | 0.26 | 0.024 (0.004) | 4.8×10^{-8} | 0.023 (0.004) | 5.8×10^{-8} | EUR | 0.85 | No | | |
| RNF219-AS1 | rs1819436 | 13 | 78,580,283 | C | T | 0.87 | 0.033 (0.006) | 6.3×10^{-9} | 0.033 (0.005) | 1.8×10^{-9} | TA/EUR | | No | rs944379 | 0.88 |
| FES | rs12906125 | 15 | 91,427,612 | G | A | 0.68 | 0.023 (0.004) | 1.7×10^{-8} | 0.023 (0.004) | 1.0×10^{-8} | TA/EUR | | No | rs6227 | 0.87 |
| IGF1R | rs7402982 | 15 | 99,193,269 | A | G | 0.43 | 0.023 (0.004) | 2.3×10^{-9} | 0.023 (0.004) | 1.1×10^{-9} | TA/EUR | | No | rs8028620 | 0.55 |
| GPR139 | rs1011939 | 16 | 19,992,996 | G | A | 0.30 | 0.022 (0.004) | 1.3×10^{-7} | 0.024 (0.004) | 2.7×10^{-9} | TA | | No | rs1858988 | 0.70 |
| CLDN7 | rs113086489 | 17 | 7,171,356 | T | C | 0.56 | 0.031 (0.004) | 9.1×10^{-16} | 0.030 (0.004) | 1.3×10^{-15} | TA/EUR | | No | rs5417 | 0.80 |
| SUZ12P1-CRLF3 | rs144843919 | 17 | 29,037,339 | G | A | 0.97 | 0.066 (0.012) | 1.4×10^{-8} | 0.068 (0.011) | 1.5×10^{-9} | TA/EUR | | No | rs8073965 | 0.22 |
| SP6-SP2 | rs12942207 | 17 | 45,968,294 | C | T | 0.30 | 0.022 (0.004) | 5.1×10^{-8} | 0.024 (0.004) | 3.0×10^{-9} | TA | | No | rs11079800 | 0.91 |
| | rs72833480 | 17 | 45,964,861 | A | G | 0.29 | 0.023 (0.004) | 4.6×10^{-8} | 0.023 (0.004) | 2.1×10^{-8} | EUR | 0.94 | No | | |
| ACTL9 | rs61154119 | 19 | 8,787,750 | T | G | 0.84 | 0.028 (0.005) | 1.1×10^{-7} | 0.028 (0.005) | 2.3×10^{-8} | TA | | No | rs2967684 | 1 |
| PEPD | rs10402712 | 19 | 33,926,013 | A | G | 0.26 | 0.022 (0.004) | 4.4×10^{-7} | 0.023 (0.004) | 2.3×10^{-8} | TA | | No | rs17833935 | 0.54 |

| Locus | SNP | Chr | Position | Alleles | | EAF | European ancestry | | Trans-ancestry | | Status ^a | EUR r^2 with lead TA | Present in HapMap? | Best proxy in HapMap | |
|-----------|------------|-----|-------------|---------|-------|------|-------------------|-----------------------|----------------|-----------------------|---------------------|------------------------|--------------------|----------------------|-----------|
| | | | (b37, bp) | Effect | Other | | β (SE) | P-value | β (SE) | P-value | | | | SNP | EUR r^2 |
| JAG1 | rs6040076 | 20 | 10,658,882 | C | G | 0.49 | 0.023 (0.004) | 2.0×10^{-9} | 0.022 (0.004) | 7.2×10^{-9} | TA/EUR | | No | rs2206815 | 0.85 |
| C20orf203 | rs28530618 | 20 | 31,275,581 | A | G | 0.49 | 0.026 (0.004) | 7.7×10^{-12} | 0.024 (0.004) | 8.4×10^{-11} | TA/EUR | | No | rs6057610 | 0.82 |
| MAFB | rs6016377 | 20 | 39,172,728 | T | C | 0.43 | 0.024 (0.004) | 9.5×10^{-10} | 0.024 (0.004) | 3.7×10^{-10} | TA/EUR | | Yes | | |
| NRIP1 | rs2229742 | 21 | 16,339,172 | G | C | 0.87 | 0.036 (0.006) | 2.2×10^{-9} | 0.034 (0.006) | 1.5×10^{-8} | TA/EUR | | Yes | | |
| KREMEN1 | rs134594 | 22 | 29,468,456 | C | T | 0.35 | 0.023 (0.004) | 1.0×10^{-8} | 0.022 (0.004) | 2.2×10^{-8} | TA/EUR | | Yes | | |
| SREBF2 | rs62240962 | 22 | 42,259,524 | C | T | 0.91 | 0.047 (0.007) | 9.7×10^{-12} | 0.047 (0.007) | 3.7×10^{-12} | TA/EUR | | No | rs10483213 | 0.70 |
| PLAC1 | rs11096402 | X | 133,827,868 | G | A | 0.28 | 0.028 (0.005) | 1.3×10^{-9} | N/A | N/A | EUR | | Yes | | |

Chr, chromosome; bp, base pair; EAF, effect allele frequency; SE, standard error.

^aStatus indicates whether the SNP was lead in European ancestry (EUR) or trans-ancestry (TA) meta-analysis. Previous denotes previously reported lead SNP⁵ at established loci.

Supplementary Table 5. Loci with multiple distinct association signals attaining genome-wide significance ($P < 5 \times 10^{-8}$) in approximate conditional meta-analysis of 143,677 individuals of European ancestry, using 5,000 white British participants from UK BioBank as a reference for linkage disequilibrium.

| Locus | Index variant | Chr | Position (b37, bp) | Alleles | EAF | Unconditional meta-analysis | | Conditional meta-analysis | | | UKBB reference r^2 | Present in HapMap ? | Best proxy in HapMap | |
|---------------|---------------|-----|--------------------|---------------|------|-----------------------------|-----------------------|---------------------------|---------------|-----------------------|----------------------|---------------------|----------------------|-----------|
| | | | | Effect/ Other | | β (SE) | P -value | Conditioned on | β (SE) | P -value | | | SNP | EUR r^2 |
| <i>ZBTB7B</i> | rs3753639 | 1 | 154,986,091 | C/T | 0.24 | 0.031 (0.004) | 7.3×10^{-12} | rs4330912 | 0.034 (0.005) | 2.2×10^{-14} | 0.019 | No | rs905938 | 0.77 |
| | rs4330912 | 1 | 155,969,428 | G/C | 0.64 | 0.021 (0.004) | 1.8×10^{-7} | rs3753639 | 0.025 (0.004) | 5.4×10^{-10} | | No | rs12043212 | 0.93 |
| <i>HMGA1</i> | rs9368777 | 6 | 33,788,637 | C/G | 0.58 | 0.022 (0.004) | 2.2×10^{-8} | rs1187118 | 0.021 (0.004) | 3.3×10^{-8} | 0.000 | No | rs4304152 | 0.57 |
| | rs1187118 | 6 | 34,169,020 | A/T | 0.17 | 0.030 (0.005) | 3.6×10^{-9} | rs9368777 | 0.030 (0.005) | 5.4×10^{-9} | | No | rs1776877 | 0.95 |
| <i>PTCH1</i> | rs12551019 | 9 | 96,949,079 | C/T | 0.68 | 0.022 (0.004) | 3.1×10^{-8} | rs3780573 | 0.024 (0.004) | 5.1×10^{-9} | 0.001 | Yes | | |
| | rs3780573 | 9 | 98,239,503 | A/G | 0.10 | 0.055 (0.006) | 7.0×10^{-18} | rs12551019 | 0.057 (0.006) | 1.2×10^{-18} | | No | rs3824488 | 1.00 |

Chr, chromosome; bp, base pair; EAF, effect allele frequency; SE, standard error.

Supplementary Table 6. Candidate gene(s) at birth weight loci.

| Locus | Lead SNP | Chr | Position (b37) | Candidate gene(s) in the locus ^a | Reports on eQTL ^b | Literature search for nearby genes (300kb) | Coding variants (EUR r^2 with the lead SNP) |
|--------------------|------------|-----|----------------|--|---|--|---|
| <i>WNT4-ZBTB40</i> | rs2473248 | 1 | 22,536,643 | <i>WNT4</i> [P,N], <i>CDC42</i> [B] | <i>WNT4</i> : Cerebellum and temporal cortex eQTL; <i>CDC42</i> : Whole blood eQTL (proxy SNP) | | - |
| <i>ZBTB7B</i> | rs3753639 | 1 | 154,986,091 | <i>SHC1</i> [B], <i>S100A7</i> [P] | | <i>SHC1</i> : The Shc adaptor proteins are key transducers of growth promotion and gene expression, and are phosphorylated by all known receptor tyrosine. | - |
| <i>FCGR2B</i> | rs72480273 | 1 | 161,644,871 | <i>ATF6</i> [B], <i>APOA2</i> [P], <i>FCGR2B</i> [N] | <i>ATF6</i> : Cerebellum and temporal cortex eQTL; <i>FCGR2B</i> : Whole blood, cerebellum and temporal cortex eQTL | <i>HSPA6</i> : higher expression of HSPA6 in placental vascular disease (PVD) vs controls; HSPA6 encodes HSP70 which was found to be upregulated in PVD vs control placenta and microvascular endothelial cells. HSP70 mRNA and protein expression also correlated negatively with infant birth weight (PMID:18372927). | - |
| <i>DTL</i> | rs61830764 | 1 | 212,289,976 | <i>DTL</i> [N] | | | - |
| <i>ATAD2B</i> | rs7575873 | 2 | 23,962,647 | <i>ATAD2B</i> [N], <i>FKBP1B</i> [P] | <i>PFN4</i> : Lymphoblastoid eQTL | | - |
| <i>EPAS1</i> | rs1374204 | 2 | 46,484,205 | <i>RHOQ</i> [B], <i>SOC5</i> [P] | <i>EPAS1</i> , <i>PRKCE</i> : Cerebellum and temporal cortex eQTL | <i>EPAS1</i> : involved in the hypoxic response and is suggested to be responsible for the genetic adaptation of high-altitude hypoxia in Tibetans. PMID 25501874 found associations of 2 variants with BW in small sample-sized Tibetans. | - |
| <i>PTH1R</i> | rs2242116 | 3 | 46,941,116 | <i>PTH1R</i> [B,N], <i>CCR1</i> [P] | <i>CCDC12</i> : Whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>ADCY5</i> | rs11719201 | 3 | 123,068,744 | <i>ADCY5</i> [B,N], <i>HCLS1</i> [P] | <i>ADCY5</i> : Cerebellum and temporal cortex eQTL (proxy SNP) | | - |
| <i>CPA3</i> | rs10935733 | 3 | 148,622,968 | <i>AGTR1</i> [B,P] | <i>CPA3</i> , <i>CPB1</i> : eQTL (proxy SNP) | <i>AGTR1</i> = angiotensin II receptor gene: good candidate as the protein mediates processes important for placentation. Levels of protein higher in umbilical cord and maternal peripheral blood in preeclampsia than in controls; levels of mRNA higher in placenta of preeclampsia than control pregnancies; birth weight negatively correlated with levels in cord (PMID 23302726). | - |
| <i>CCNL1-LEKR1</i> | rs13322435 | 3 | 156,795,468 | <i>LEKR1</i> [N], <i>IL12A</i> [P] | <i>LEKR1</i> : Temporal cortex eQTL; <i>CCNL1</i> : Whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>LCORL</i> | rs925098 | 4 | 17,919,811 | <i>LCORL</i> [N], <i>SLIT2</i> [P] | <i>LCORL</i> : Cerebellum and temporal cortex eQTL | Adipose tissue and muscle expression in cattle of <i>NCAPG</i> and <i>LCORL</i> associated with feed intake and weight gain in cattle (these are in a QTL for body weight in cattle). | - |
| <i>HHIP</i> | rs6537307 | 4 | 145,601,863 | <i>HHIP</i> [N], <i>ANAPC10</i> [P] | <i>HHIP</i> : Cerebellum and temporal cortex eQTL | | - |
| <i>5q11.2</i> | rs854037 | 5 | 57,091,783 | <i>ACTBL2</i> [N], <i>IL6ST</i> [P] | | | - |
| <i>EBF1</i> | rs7729301 | 5 | 157,886,953 | <i>EBF1</i> [N,B], <i>FABP6</i> [P] | | | - |
| <i>CDKAL1</i> | rs35261542 | 6 | 20,675,792 | <i>CDKAL1</i> [B], <i>SOX4</i> [P] | <i>CDKAL1</i> : Cerebellum and temporal cortex eQTL | | - |

| Locus | Lead SNP | Chr | Position (b37) | Candidate gene(s) in the locus ^a | Reports on eQTL ^b | Literature search for nearby genes (300kb) | Coding variants (EUR r^2 with the lead SNP) |
|--------------------|-------------|-----|----------------|--|--|---|---|
| <i>HIST1H2BE</i> | rs9379832 | 6 | 26,186,200 | <i>HIST1H2BE</i> [N,B], <i>BTN2A2</i> [P] | | | - |
| <i>HMGA1</i> | rs7742369 | 6 | 34,165,721 | <i>HMGA1</i> [N,B], <i>GRM4</i> [B], <i>MAPK14</i> [P] | <i>HMGA1</i> : Cerebellum and temporal cortex eQTL | | - |
| <i>L3MBTL3</i> | rs1415701 | 6 | 130,345,835 | <i>L3MBTL3</i> [E,N], <i>SAMD3</i> [E], <i>ENPP1</i> [P] | <i>L3MBTL3</i> , <i>SAMD3</i> : Whole blood, cerebellum and temporal cortex eQTLs | | - |
| <i>ESR1</i> | rs1101081 | 6 | 152,032,917 | <i>ESR1</i> [N,B], <i>LATS1</i> [P] | <i>ESR1</i> : Monocyte, cerebellum and temporal cortex eQTL (for proxy SNP) | | - |
| <i>GNA12</i> | rs798489 | 7 | 2,801,803 | <i>GNA12</i> [N,E], <i>PDGFA</i> [N] | <i>GNA12</i> : Nerve tibial, lymphoblastid, whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>IGF2BP3</i> | rs11765649 | 7 | 23,479,013 | <i>IGF2BP3</i> [N,B], <i>NPY</i> [P] | <i>CCDC126</i> : lymphoblastoid eQTL | | - |
| <i>TBX20</i> | rs6959887 | 7 | 35,295,365 | <i>TBX20</i> [N], <i>SEPT7</i> [P] | <i>TBX20</i> : Cerebellum and temporal cortex eQTL | | - |
| <i>YKT6-GCK</i> | rs138715366 | 7 | 44,246,271 | <i>YKT6</i> [N], <i>GCK</i> [B] | | GCK mutations reduce birth weight and increase fasting glucose (Hattersley et al 1999, Nat Genet). | - |
| <i>MLXIPL</i> | rs62466330 | 7 | 73,056,805 | <i>MLXIPL</i> [N], <i>FZD9</i> [P] | <i>MLXIPL</i> : Cerebellum and temporal cortex eQTL (for proxy SNP) | | - |
| <i>ANK1-NKX6-3</i> | rs13266210 | 8 | 41,533,514 | <i>ANK1</i> [N,B], <i>SFRP1</i> [P] | <i>ANK1</i> : Cerebellum, temporal cortex and lung eQTL | | - |
| <i>TRIB1</i> | rs6989280 | 8 | 126,508,746 | <i>TRIB1</i> [N], <i>MYC</i> [P] | <i>TRIB1</i> : Cerebellum and temporal cortex eQTL | | - |
| <i>SLC45A4</i> | rs12543725 | 8 | 142,247,979 | <i>SLC45A4</i> [N,E], <i>PTK2</i> [B,P] | <i>SLC45A4</i> : Whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>PTCH1</i> | rs28510415 | 9 | 98,245,026 | <i>PTCH1</i> [N,B], <i>FOXE1</i> [P] | <i>PTCH1</i> : Whole blood, cerebellum and temporal cortex eQTL (for proxy SNP) | <i>FANCC</i> : mutations result in Fanconi anaemia (associated with low birth weight and shorter stature). | - |
| <i>LPAR1</i> | rs2150052 | 9 | 113,945,067 | <i>LPAR1</i> [N,E,B,P] | <i>LPAR1</i> : whole blood eQTL | | - |
| <i>PHF19</i> | rs7847628 | 9 | 123,631,225 | <i>C5</i> [B], <i>DAB2IP</i> [P] | <i>C5</i> : naïve monocyte, whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>STRBP</i> | rs700059 | 9 | 125,824,055 | <i>STRBP</i> [E], <i>C5</i> [P] | <i>STRBP</i> : Whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>HHEX-IDE</i> | rs61862780 | 10 | 94,468,643 | <i>IDE</i> [B], <i>PLCE1</i> [P] | <i>HHEX</i> : monocyte, whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>NTSC2</i> | rs74233809 | 10 | 104,913,940 | <i>NTSC2</i> [N], <i>CHUK</i> [P] | | <i>CYP17A1</i> : mutation in maternal <i>CYP17A1</i> is associated with small for gestational age (SGA) (PMID 14665706) | - |
| <i>ADRB1</i> | rs7076938 | 10 | 115,789,375 | <i>ADRB1</i> [N,C,B], <i>AFAP1L2</i> [P] | <i>ADRB1</i> : cerebellum and temporal cortex eQTL | | rs1801253 (G389R, $r^2=0.99$, damaging) |
| <i>PLEKHA1</i> | rs2421016 | 10 | 124,167,512 | <i>PLEKHA1</i> [N,E], <i>FGFR2</i> [P] | <i>PLEKHA1</i> : Whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>INS-IGF2</i> | rs72851023 | 11 | 2,130,620 | <i>INS-IGF2</i> [N,B,imp] | | | - |

| Locus | Lead SNP | Chr | Position (b37) | Candidate gene(s) in the locus ^a | Reports on eQTL ^b | Literature search for nearby genes (300kb) | Coding variants (EUR r^2 with the lead SNP) |
|----------------------|-------------|-----|----------------|--|---|--|---|
| <i>MTNR1B</i> | rs10830963 | 11 | 92,708,710 | <i>MTNR1B</i> [N,B,P] | <i>MTNR1B</i> : Cerebellum and temporal cortex eQTL | | - |
| <i>APOLD1</i> | rs11055034 | 12 | 12,890,626 | <i>CDKN1B</i> [B,E,P], <i>HEBP1</i> [B] | <i>CDKN1B</i> : Cerebellum and temporal cortex eQTL | | - |
| <i>ABCC9</i> | rs139975827 | 12 | 22,068,161 | <i>ABCC9</i> [N], <i>KRAS</i> [P] | | | - |
| <i>ITPR2</i> | rs12823128 | 12 | 26,872,730 | <i>ITPR2</i> [N,B] | <i>ITPR2</i> : Cerebellum and temporal cortex eQTL | | - |
| <i>HMGA2</i> | rs1351394 | 12 | 66,351,826 | <i>HMGA2</i> [N,B], <i>IRAK3</i> [P] | <i>HMGA2</i> : Cerebellum and temporal cortex eQTL | <i>HMGA2</i> : 12q14 microdeletion syndrome implicates <i>HMGA2</i> as causal to height (PMID: 17220210 and 19298872). | - |
| <i>IGF1</i> | rs7964361 | 12 | 102,994,878 | <i>IGF1</i> [N,B,P] | | | - |
| <i>LINC00332</i> | rs2324499 | 13 | 40,662,001 | <i>LINC00332</i> [N], <i>TNFSF11</i> [P] | | | - |
| <i>RB1</i> | rs2854355 | 13 | 48,882,363 | <i>RB1</i> [N,B,P,imp], <i>LPAR6</i> [imp,B] | <i>RB1</i> : Whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>RNF219-AS1</i> | rs1819436 | 13 | 78,580,283 | <i>EDNRB</i> [B,P] | <i>EDNRB</i> : Monocyte, cerebellum and temporal cortex eQTL (for proxy SNP) | | - |
| <i>FES</i> | rs12906125 | 15 | 91,427,612 | <i>FES</i> [N,E], <i>MESP1</i> [P] | <i>FES</i> : Thyroid, whole blood, cerebellum, temporal cortex and tibial nerve eQTL | <i>FURIN</i> : a ubiquitous proprotein convertase, the major processing enzyme of the secretory pathway. It is involved in processing proinsulin to insulin, and in processing proIGF2 (PMID 9660813). | - |
| <i>IGF1R</i> | rs7402982 | 15 | 99,193,269 | <i>IGF1R</i> [B] | | | - |
| <i>GPR139</i> | rs1011939 | 16 | 19,992,996 | <i>GPR139</i> [N], <i>RPS15A</i> [P] | | | - |
| <i>CLDN7</i> | rs113086489 | 17 | 7,171,356 | <i>SLC2A4</i> [B], <i>POLR2A</i> [S], <i>DVL2</i> [P] | <i>SLC2A4</i> : Cerebellum and temporal cortex eQTL; <i>DVL2</i> : monocyte, whole blood, cerebellum and temporal cortex eQTL (for proxy SNP) | <i>SLC2A4</i> (<i>GLUT4</i>): an insulin responsive glucose transporter. PMID 24062248 showed lower mRNA in preadipocytes from LBW vs NBW. Protein level of <i>SLC2A4</i> was lower in adipose tissue of LBW men compared to NMW (PMID 17063325). | - |
| <i>SUZ12P1-CRLF3</i> | rs144843919 | 17 | 29,037,339 | <i>SUZ12P1</i> [N] | | | - |
| <i>SP6-SP2</i> | rs12942207 | 17 | 45,968,294 | <i>SP2</i> [N], <i>NGFR</i> [P] | <i>SP2</i> : Whole blood, cerebellum and temporal cortex eQTL (for proxy SNP) | | - |
| <i>ACTL9</i> | rs61154119 | 19 | 8,787,750 | <i>ACTL9</i> [N], <i>INSR</i> [P] | | | - |
| <i>PEPD</i> | rs10402712 | 19 | 33,926,013 | <i>CEBPA</i> [B] | | <i>CEBPA</i> : an adipogenic transcription factor involved in adipocyte differentiation. Expression levels lower in smaller pig fetuses (PMID 21354690). | - |
| <i>JAG1</i> | rs6040076 | 20 | 10,658,882 | <i>JAG1</i> [N], <i>MKKS</i> [P] | <i>JAG1</i> : Cerebellum and temporal cortex eQTL (proxy SNP) | <i>JAG1</i> is a ligand that interacts with receptors in the notch signalling pathway. Notch pathway plays an important role during placental development; notch pathway downregulation is associated with preeclampsia, and specifically <i>JAG1</i> mRNA levels downregulated in preeclampsia placental samples vs controls (PMID 25962154). | - |
| <i>C20orf203</i> | rs28530618 | 20 | 31,275,581 | <i>BAK1P1</i> [E], <i>FOXS1</i> [P] | <i>BAK1P1</i> : thyroid eQTL (for proxy SNP) | | - |

| Locus | Lead SNP | Chr | Position (b37) | Candidate gene(s) in the locus ^a | Reports on eQTL ^b | Literature search for nearby genes (300kb) | Coding variants (EUR r^2 with the lead SNP) |
|----------------|------------|-----|----------------|---|--|--|---|
| <i>MAFB</i> | rs6016377 | 20 | 39,172,728 | <i>MAFB</i> [N], <i>PLCG1</i> [P] | <i>MAFB</i> : Cerebellum and temporal cortex eQTL (for proxy SNP) | Lots of literature on importance of <i>MAFB</i> in transcriptional regulation in embryonic islets, e.g. analysis of <i>Mafa</i> (-/-) and pancreas-specific <i>Mafa</i> (Δ panc) deletion mutant mice demonstrated a primary role for <i>MafA</i> in adult β -cell activity, different from the embryonic importance of <i>MafB</i> (PMID 24520122). | - |
| <i>NRIP1</i> | rs2229742 | 21 | 16,339,172 | <i>NRIP1</i> [N,C], <i>SAMSN1</i> [P] | <i>NRIP1</i> : Cerebellum and temporal cortex eQTL | | rs2229742 (R448G, lead SNP, damaging) |
| <i>KREMEN1</i> | rs134594 | 22 | 29,468,456 | <i>KREMEN1</i> [N,E], <i>YWHAH</i> [P] | <i>KREMEN1</i> : Monocyte, heart, whole blood, cerebellum and temporal cortex eQTL | | - |
| <i>SREBF2</i> | rs62240962 | 22 | 42,259,524 | <i>SREBF2</i> [N,B], <i>PDGFB</i> [P] | | | - |
| <i>PLAC1</i> | rs11096402 | X | 133,827,868 | <i>PLAC1</i> [N] | | <i>PLAC1</i> shows placenta-specific expression in human (PMID: 10995572). A mutant mouse model showed <i>PLAC1</i> as a paternally imprinted, X-linked gene essential for normal placental and embryonic development (PMID: 22729990). | - |

^aB = biological candidate, C = coding variant, E = eQTL (from GTEx), N=nearest, P = best ranked gene in PPI network, imp = imprinted gene.

^beQTL reported from GTEx v4, GEUVADIS, and 11 other studies for the lead variant. When there was no eQTL report for the lead variant, eQTL reports for the proxy SNPs ($r^2 > 0.8$ with the lead variant) are shown.

Supplementary Table 7. Summary of 99% credible sets at 62 distinct autosomal association signals.

| Locus | Index variant | Chr | Position (b37, bp) | MANTRA log ₁₀ BF | 99% credible set | | |
|--------------------|---------------|-----|-----------------------|--------------------------------|------------------|---------------|-------------------------|
| | | | | | Variants | Distance (bp) | Interval (bp) |
| <i>WNT4-ZBTB40</i> | rs2473248 | 1 | 22,536,643 | 7.68 | 7 | 17,534 | 22,536,643-22,554,176 |
| <i>ZBTB7B</i> | rs3753639 | 1 | 154,986,091 | 12.19 | 6 | 22,088 | 154,984,363-155,006,450 |
| | rs4330912 | 1 | 155,969,428 | 7.93 | 45 | 403,839 | 155,612,197-156,016,035 |
| <i>FCGR2B</i> | rs72480273 | 1 | 161,644,871 | 7.48 | 37 | 173,221 | 161,500,712-161,673,932 |
| <i>DTL</i> | rs61830764 | 1 | 212,289,976 | 6.02 | 56 | 702,234 | 211,830,044-212,532,277 |
| <i>ATAD2B</i> | rs7575873 | 2 | 23,962,647 | 8.87 | 234 | 477,824 | 23,887,437-24,365,260 |
| <i>EPAS1</i> | rs1374204 | 2 | 46,484,205 | 27.22 | 4 | 695 | 46,484,188-46,484,882 |
| <i>PTH1R</i> | rs2242116 | 3 | 46,941,116 | 6.65 | 54 | 243,837 | 46,925,539-47,169,375 |
| <i>ADCY5</i> | rs11719201 | 3 | 123,068,744 | 24.38 | 6 | 16,621 | 123,065,778-123,082,398 |
| <i>CPA3</i> | rs10935733 | 3 | 148,622,968 | 7.91 | 80 | 33,654 | 148,591,725-148,625,378 |
| <i>CCNL1-LEKR1</i> | rs13322435 | 3 | 156,795,468 | 40.24 | 11 | 3,265 | 156,795,468-156,798,732 |
| <i>LCORL</i> | rs925098 | 4 | 17,919,811 | 13.56 | 15 | 123,806 | 17,901,679-18,025,484 |
| <i>HHIP</i> | rs6537307 | 4 | 145,601,863 | 10.43 | 43 | 137,679 | 145,520,608-145,658,286 |
| <i>5q11.2</i> | rs854037 | 5 | 57,091,783 | 5.77 | 63 | 276,269 | 56,906,160-57,182,428 |
| <i>EBF1</i> | rs7729301 | 5 | 157,886,953 | 7.62 | 50 | 77380 | 157,884,706-157,962,085 |
| <i>CDKAL1</i> | rs35261542 | 6 | 20,675,792 | 26.41 | 7 | 14,242 | 20,673,880-20,688,121 |
| <i>HIST1H2BE</i> | rs9379832 | 6 | 26,186,200 | 6.64 | 5 | 26,542 | 26,180,634-26,207,175 |
| <i>HMGA1</i> | rs9368777 | 6 | 33,788,637 | 6.20 | 91 | 570,460 | 33,292,543-33,863,002 |
| | rs1187118 | 6 | 34,169,020 | 6.94 | 27 | 148,294 | 34,095,696-34,243,989 |
| <i>L3MBTL3</i> | rs1415701 | 6 | 130,345,835 | 9.03 | 2 | 4,601 | 130,341,235-130,345,835 |
| <i>ESR1</i> | rs1101081 | 6 | 152,032,917 | 17.69 | 14 | 20,968 | 152,029,556-152,050,523 |
| <i>GNA12</i> | rs798489 | 7 | 2,801,803 | 7.01 | 114 | 160,777 | 2,752,152-2,912,928 |
| <i>IGF2BP3</i> | rs11765649 | 7 | 23,479,013 | 7.58 | 23 | 111,149 | 23,474,687-23,585,835 |
| <i>TBX20</i> | rs6959887 | 7 | 35,295,365 | 6.75 | 113 | 109,117 | 35,196,208-35,305,324 |
| <i>YKT6-GCK</i> | rs138715366 | 7 | 44,246,271 | 24.10 | 1 | 1 | 44,246,271-44,246,271 |
| <i>MLXIPL</i> | rs62466330 | 7 | 73,056,805 | 9.60 | 102 | 221,896 | 72,836,230-73,058,025 |
| <i>ANK1-NKX6-3</i> | rs13266210 | 8 | 41,533,514 | 9.22 | 17 | 28,742 | 41,508,577-41,537,318 |
| <i>TRIB1</i> | rs6989280 | 8 | 126,508,746 | 5.69 | 39 | 411,117 | 126,114,947-126,526,063 |
| <i>SLC45A4</i> | rs12543725 | 8 | 142,247,979 | 7.18 | 10 | 13,200 | 142,239,381-142,252,580 |
| <i>PTCH1</i> | rs12551019 | 9 | 96,949,079 | 6.96 | 40 | 652,934 | 96,900,505-97,553,438 |
| | rs3780573 | 9 | 98,239,503 | 16.44 | 23 | 53,294 | 98,212,608-98,265,901 |

| Locus | Index variant | Chr | Position | MANTRA log ₁₀ BF | 99% credible set | | |
|----------------------|---------------|-----|-------------|--------------------------------|------------------|---------------|-------------------------|
| | | | (b37, bp) | | Variants | Distance (bp) | Interval (bp) |
| <i>LPAR1</i> | rs2150052 | 9 | 113,945,067 | 6.29 | 60 | 214,738 | 113,828,811-114,043,548 |
| <i>PHF19</i> | rs7847628 | 9 | 123,631,225 | 6.95 | 5 | 90,924 | 123,631,225-123,722,148 |
| <i>STRBP</i> | rs700059 | 9 | 125,824,055 | 10.35 | 135 | 432,744 | 125,605,840-126,038,583 |
| <i>HHEX-IDE</i> | rs61862780 | 10 | 94,468,643 | 12.40 | 31 | 134,694 | 94,358,023-94,492,716 |
| <i>NT5C2</i> | rs74233809 | 10 | 104,913,940 | 7.25 | 98 | 404,547 | 104,655,350-105,059,896 |
| <i>ADRB1</i> | rs7076938 | 10 | 115,789,375 | 15.69 | 4 | 9,521 | 115,789,375-115,798,895 |
| <i>PLEKHA1</i> | rs2421016 | 10 | 124,167,512 | 6.90 | 26 | 70,596 | 124,127,990-124,198,585 |
| <i>INS-IGF2</i> | rs72851023 | 11 | 2,130,620 | 7.89 | 48 | 145,434 | 2,078,742-2,224,175 |
| <i>MTNR1B</i> | rs10830963 | 11 | 92,708,710 | 5.78 | 52 | 499,464 | 92,225,858-92,725,321 |
| <i>APOLD1</i> | rs11055034 | 12 | 12,890,626 | 6.33 | 13 | 42,180 | 12,849,241-12,891,420 |
| <i>ABCC9</i> | rs139975827 | 12 | 22,068,161 | 5.64 | 72 | 133,773 | 22,027,182-22,160,954 |
| <i>ITPR2</i> | rs12823128 | 12 | 26,872,730 | 6.12 | 114 | 291,548 | 26,759,026-27,050,573 |
| <i>HMGA2</i> | rs1351394 | 12 | 66,351,826 | 30.74 | 7 | 32,282 | 66,343,810-66,376,091 |
| <i>IGF1</i> | rs7964361 | 12 | 102,994,878 | 6.71 | 42 | 213,557 | 102,867,636-103,081,192 |
| <i>LINC00332</i> | rs2324499 | 13 | 40,662,001 | 6.83 | 41 | 95,318 | 40,583,503-40,678,820 |
| <i>RB1</i> | rs2854355 | 13 | 48,882,363 | 6.23 | 143 | 403,044 | 48,797,661-49,200,704 |
| <i>RNF219-AS1</i> | rs1819436 | 13 | 78,580,283 | 7.44 | 105 | 220,228 | 78,443,297-78,663,524 |
| <i>FES</i> | rs12906125 | 15 | 91,427,612 | 6.68 | 24 | 36,382 | 91,404,705-91,441,086 |
| <i>IGF1R</i> | rs7402982 | 15 | 99,193,269 | 7.59 | 5 | 15,682 | 99,177,595-99,193,276 |
| <i>GPR139</i> | rs1011939 | 16 | 19,992,996 | 7.20 | 55 | 53,120 | 19,992,996-20,461,115 |
| <i>CLDN7</i> | rs113086489 | 17 | 7,171,356 | 13.41 | 29 | 84,570 | 7,101,292-7,185,861 |
| <i>SUZ12P1-CRLF3</i> | rs144843919 | 17 | 29,037,339 | 7.60 | 21 | 520,396 | 28,780,178-29,300,573 |
| <i>SP6-SP2</i> | rs12942207 | 17 | 45,968,294 | 7.04 | 21 | 90,685 | 45,940,310-46,030,994 |
| <i>ACTL9</i> | rs61154119 | 19 | 8,787,750 | 6.24 | 21 | 4,455 | 8,785,744-8,790,198 |
| <i>PEPD</i> | rs10402712 | 19 | 33,926,013 | 6.40 | 35 | 218,599 | 33,784,657-34,003,255 |
| <i>JAG1</i> | rs6040076 | 20 | 10,658,882 | 6.72 | 43 | 164,577 | 10,539,775-10,704,351 |
| <i>C20orf203</i> | rs28530618 | 20 | 31,275,581 | 8.69 | 63 | 114,192 | 31,214,234-31,328,425 |
| <i>MAFB</i> | rs6016377 | 20 | 39,172,728 | 8.15 | 33 | 72,001 | 39,144,286-39,216,286 |
| <i>NRIP1</i> | rs2229742 | 21 | 16,339,172 | 6.69 | 4 | 85,250 | 16,339,172-16,424,421 |
| <i>KREMEN1</i> | rs134594 | 22 | 29,468,456 | 6.41 | 56 | 52,177 | 29,461,150-29,513,326 |
| <i>SREBF2</i> | rs62240962 | 22 | 42,259,524 | 10.01 | 4 | 189,151 | 42,070,374-42,259,524 |

Chr, chromosome; bp, base pair; BF, Bayes factor.

Supplementary Table 8. BW credible set enrichment for DNaseI hypersensitive sites.

(a) 128 cell-type DHS annotation and 4 genic annotation tested for enrichment individually

| Cell type | Lower 95%CI ^a | Effect ^a | Upper 95%CI ^a | Tissue | Category |
|------------------------|--------------------------|---------------------|--------------------------|-----------------------|-----------------|
| iPS_6_9 | -20.00 | 0.35 | 1.74 | Stem cell | Stem cell |
| iPS_4_7 | -20.00 | 0.82 | 2.04 | Stem cell | Stem cell |
| iPS_19_7 | -1.22 | 0.96 | 2.09 | Stem cell | Stem cell |
| iPS_19_11 | -1.40 | 0.80 | 1.93 | Stem cell | Stem cell |
| hESCT0 | -0.10 | 1.26 | 2.16 | Stem cell | Stem cell |
| HESC | -20.00 | 0.21 | 1.79 | Stem cell | Stem cell |
| NPC | 0.94 | 2.14 | 2.99 | Brain | Progenitor cell |
| HMVEC_dNeo | 0.24 | 1.62 | 2.54 | Neonatal blood vessel | Neonatal |
| HMVEC_dLyNeo | 0.48 | 1.71 | 2.58 | Neonatal blood vessel | Neonatal |
| HMVEC_dBiNeo | -0.82 | 1.00 | 2.01 | Neonatal blood vessel | Neonatal |
| NHDF_Neo | 1.22 | 2.29 | 3.12 | Skin | Neonatal |
| fThymus | 0.88 | 1.94 | 2.75 | Foetal thymus | Foetal |
| fTestes | 0.88 | 1.97 | 2.79 | Foetal testes | Foetal |
| fStomach | 0.40 | 1.63 | 2.50 | Foetal stomach | Foetal |
| fSpleen | 0.75 | 1.86 | 2.68 | Foetal spleen | Foetal |
| fSpinal_cord | 0.52 | 1.58 | 2.38 | Foetal skin | Foetal |
| fSkin_fibro_upper_back | -0.86 | 1.43 | 2.44 | Foetal skin | Foetal |
| fSkin_fibro_scalp | -4.83 | 1.03 | 2.17 | Foetal skin | Foetal |
| fSkin_fibro_leg_R_quad | -20.00 | 0.84 | 2.11 | Foetal skin | Foetal |
| fSkin_fibro_leg_L_quad | -0.89 | 1.23 | 2.25 | Foetal skin | Foetal |
| fSkin_fibro_bicep_R | -20.00 | 0.73 | 2.05 | Foetal skin | Foetal |
| fSkin_fibro_bicep_L | -20.00 | 1.09 | 2.24 | Foetal skin | Foetal |
| fSkin_fibro_back | -20.00 | 0.04 | 1.75 | Foetal skin | Foetal |
| fSkin_fibro_abdomen | -3.50 | 1.30 | 2.41 | Foetal skin | Foetal |
| fSkin | -20.00 | 1.13 | 2.31 | Foetal skin | Foetal |
| fPlacenta | 0.80 | 1.72 | 2.49 | Foetal placenta | Foetal |
| fMuscle_upper_trunk | 0.70 | 1.74 | 2.55 | Foetal muscle | Foetal |
| fMuscle_upper_limb_sk | -0.08 | 1.43 | 2.35 | Foetal muscle | Foetal |
| fMuscle_upper_back | 0.48 | 1.70 | 2.54 | Foetal muscle | Foetal |
| fMuscle_trunk | 0.63 | 1.72 | 2.54 | Foetal muscle | Foetal |
| fMuscle_lower_limb | 0.53 | 1.53 | 2.33 | Foetal muscle | Foetal |
| fMuscle_leg | 0.22 | 1.60 | 2.49 | Foetal muscle | Foetal |
| fMuscle_back | 0.34 | 1.37 | 2.17 | Foetal muscle | Foetal |
| fMuscle_arm | 0.46 | 1.73 | 2.60 | Foetal muscle | Foetal |
| fLung_R | 0.76 | 1.84 | 2.65 | Foetal lung | Foetal |
| fLung_L | 0.09 | 1.50 | 2.41 | Foetal lung | Foetal |
| fLung | -1.03 | 1.34 | 2.42 | Foetal lung | Foetal |
| fKidney_renal_pelvis_R | 1.23 | 2.12 | 2.87 | Foetal kidney | Foetal |
| fKidney_renal_pelvis_L | 0.51 | 1.62 | 2.45 | Foetal kidney | Foetal |
| fKidney_renal_pelvis | 0.60 | 1.69 | 2.52 | Foetal kidney | Foetal |
| fKidney_renal_cortex_R | 0.03 | 1.44 | 2.35 | Foetal kidney | Foetal |
| fKidney_renal_cortex_L | 0.91 | 1.88 | 2.67 | Foetal kidney | Foetal |
| fKidney_R | 0.52 | 1.63 | 2.45 | Foetal kidney | Foetal |
| fKidney_L | 0.70 | 1.81 | 2.65 | Foetal kidney | Foetal |
| fKidney | 0.42 | 1.65 | 2.51 | Foetal kidney | Foetal |
| fIntestine_Sm | -0.17 | 1.29 | 2.22 | Foetal intestine | Foetal |
| fIntestine_Lg | 0.25 | 1.63 | 2.55 | Foetal intestine | Foetal |
| fHeart | 0.21 | 1.56 | 2.45 | Foetal heart | Foetal |
| fBrain | 0.65 | 1.90 | 2.78 | Foetal brain | Foetal |

| Cell type | Lower 95%_CI ^a | Effect ^a | Upper 95%_CI ^a | Tissue | Category |
|--------------------|------------------------------|---------------------|------------------------------|----------------|-----------|
| fAdrenal | 0.54 | 1.69 | 2.52 | Foetal adrenal | Foetal |
| WI_38_TAM | -3.99 | 0.83 | 1.95 | Embryonic | Embryonic |
| WI_38 | -20.00 | 0.43 | 1.87 | Embryonic | Embryonic |
| Trophoblast | -20.00 | 1.21 | 2.36 | Embryonic | Embryonic |
| Mesendoderm | -0.52 | 1.15 | 2.15 | Embryonic | Embryonic |
| Skin_Melanocytes | -0.05 | 1.40 | 2.33 | Skin | Adult |
| Skin_Keratinocytes | -20.00 | 0.84 | 2.00 | Skin | Adult |
| Skin_Fibroblasts | 0.81 | 1.81 | 2.64 | Skin | Adult |
| NHEK | -20.00 | 0.57 | 1.98 | Skin | Adult |
| NHDF_Ad | -0.75 | 1.09 | 2.09 | Skin | Adult |
| BJ | -0.05 | 1.65 | 2.64 | Skin | Adult |
| AG10803 | -0.12 | 1.50 | 2.46 | Skin | Adult |
| AG09309 | -20.00 | 1.03 | 2.20 | Skin | Adult |
| AG04449 | -0.32 | 1.50 | 2.47 | Skin | Adult |
| PrEC | -20.00 | 0.83 | 2.13 | Prostate | Adult |
| LNCap | -20.00 | 1.28 | 2.42 | Prostate | Adult |
| SkMC | -0.82 | 1.33 | 2.38 | Muscle | Adult |
| HSMM_D | -20.00 | 0.66 | 1.83 | Muscle | Adult |
| HSMM | -20.00 | -1.29 | 1.31 | Muscle | Adult |
| NHLF | -2.24 | 0.96 | 2.08 | Lung | Adult |
| IMR90 | -20.00 | 0.61 | 1.93 | Lung | Adult |
| HPF | -1.69 | 1.00 | 2.10 | Lung | Adult |
| AG04450 | 0.03 | 1.60 | 2.56 | Lung | Adult |
| HRGEC | -0.64 | 1.57 | 2.60 | Kidney | Adult |
| HCM | -20.00 | 0.79 | 1.99 | Heart | Adult |
| HCFaa | -0.48 | 1.25 | 2.25 | Heart | Adult |
| HCF | -2.09 | 1.13 | 2.20 | Heart | Adult |
| HGF | 0.49 | 1.88 | 2.79 | Gingival | Adult |
| AG09319 | -0.49 | 1.54 | 2.56 | Gingival | Adult |
| HFF_MyC | -1.06 | 1.16 | 2.21 | Foreskin | Adult |
| HFF | 0.51 | 1.74 | 2.62 | Foreskin | Adult |
| HConF | -20.00 | 0.94 | 2.12 | Eye | Adult |
| SAEC | -20.00 | 0.92 | 2.11 | Epithelium | Adult |
| RPTEC | -20.00 | -0.11 | 1.55 | Epithelium | Adult |
| HRPEpiC | -20.00 | 0.20 | 1.67 | Epithelium | Adult |
| HRE | -1.34 | 1.17 | 2.20 | Epithelium | Adult |
| HRCE | -20.00 | 0.82 | 2.02 | Epithelium | Adult |
| HPdLF | -20.00 | 0.55 | 1.91 | Epithelium | Adult |
| HNPCEpiC | -0.11 | 1.42 | 2.36 | Epithelium | Adult |
| HIPEpiC | -0.16 | 1.38 | 2.32 | Epithelium | Adult |
| HEEpiC | -0.05 | 1.51 | 2.45 | Epithelium | Adult |
| HCPEpiC | 0.06 | 1.53 | 2.45 | Epithelium | Adult |
| HAepiC | -0.87 | 1.21 | 2.23 | Epithelium | Adult |
| HVMF | -0.07 | 1.53 | 2.49 | Connective | Adult |
| vHMEC | -0.13 | 1.36 | 2.29 | Breast | Adult |
| HMF | -0.37 | 1.35 | 2.31 | Breast | Adult |
| HMEC | 0.27 | 1.69 | 2.60 | Breast | Adult |
| NHA | -1.48 | 1.16 | 2.22 | Brain | Adult |
| HAsp | -20.00 | 0.63 | 1.91 | Brain | Adult |
| HAh | -20.00 | 0.59 | 1.80 | Brain | Adult |
| HAc | -20.00 | 0.63 | 1.97 | Brain | Adult |
| HUVEC | -0.57 | 1.36 | 2.38 | Blood vessel | Adult |

| Cell type | Lower 95%_CI ^a | Effect ^a | Upper 95%_CI ^a | Tissue | Category |
|---------------|---------------------------|---------------------|---------------------------|--------------|----------|
| HPAF | -20.00 | 0.95 | 2.08 | Blood vessel | Adult |
| HPAEC | -1.23 | 1.34 | 2.44 | Blood vessel | Adult |
| HMVEC_LLy | -0.39 | 1.32 | 2.34 | Blood vessel | Adult |
| HMVEC_LBI | -4.39 | 0.70 | 1.90 | Blood vessel | Adult |
| HMVEC_dLyAd | -20.00 | 1.02 | 2.24 | Blood vessel | Adult |
| HMVEC_dBIAd | 0.03 | 1.46 | 2.38 | Blood vessel | Adult |
| HMVEC_dAd | -0.79 | 1.37 | 2.44 | Blood vessel | Adult |
| HBMEC | -1.98 | 0.88 | 1.96 | Blood vessel | Adult |
| AoAF | -20.00 | 0.73 | 1.92 | Blood vessel | Adult |
| hTH2 | 0.28 | 1.54 | 2.43 | Blood | Adult |
| hTH17 | -20.00 | 1.00 | 2.39 | Blood | Adult |
| hTH1 | 0.19 | 1.61 | 2.53 | Blood | Adult |
| H9_P42 | -0.80 | 1.24 | 2.27 | Blood | Adult |
| H1_P18 | -0.07 | 1.40 | 2.35 | Blood | Adult |
| GM12878 | -0.72 | 1.24 | 2.30 | Blood | Adult |
| GM12865 | -20.00 | 0.73 | 2.07 | Blood | Adult |
| GM12864 | -20.00 | 0.85 | 2.10 | Blood | Adult |
| GM06990 | -20.00 | 0.98 | 2.19 | Blood | Adult |
| CD8 | 0.34 | 1.73 | 2.66 | Blood | Adult |
| CD56 | -1.01 | 1.06 | 2.13 | Blood | Adult |
| CD4 | -0.49 | 1.28 | 2.29 | Blood | Adult |
| CD34 | -0.15 | 1.50 | 2.49 | Blood | Adult |
| CD3_CordBlood | -1.52 | 1.56 | 2.69 | Blood | Adult |
| CD3 | -2.48 | 1.09 | 2.30 | Blood | Adult |
| CD20 | -2.52 | 0.90 | 2.01 | Blood | Adult |
| CD19 | 0.29 | 1.71 | 2.66 | Blood | Adult |
| CD14 | -0.70 | 1.07 | 2.10 | Blood | Adult |
| TSS | -0.87 | 1.37 | 2.42 | Genic | Genic |
| 3UTR | -20.00 | 1.04 | 2.47 | Genic | Genic |
| 5UTR | -47.50 | -2.00 | 2.14 | Genic | Genic |
| CDS | -2.08 | 1.03 | 2.48 | Genic | Genic |

(b) Six categorical fields from ENCODE tested for enrichment in a joint model

| Annotation | Lower 95% CI ^a | Effect ^a | Upper 95% CI ^a |
|---------------|---------------------------|---------------------|---------------------------|
| Neonatal DHS | 0.60 | 1.59 | 2.40 |
| Foetal DHS | 0.57 | 1.43 | 2.37 |
| Genic | -0.85 | 0.63 | 1.61 |
| Stem_cell DHS | -1.35 | 0.09 | 1.04 |
| Adult DHS | -1.12 | -0.26 | 0.62 |
| Embryonic DHS | -3.95 | -1.49 | -0.39 |

^aEstimated effects of enrichment and 95% CI are natural log transformed. We considered an annotation enriched if the 95% CI of the estimated effects of enrichment did not overlap zero.

Supplementary Table 9. Genetic variance explained.

| Study | <i>N</i> | By 62 known and novel SNPs | | By 55 novel SNPs | |
|--------------|----------|----------------------------|-------|--------------------|-------|
| | | Variance explained | SE | Variance explained | SE |
| HAPO | 1338 | 0.020 | 0.011 | 0.013 | 0.010 |
| NFBC1966 | 5402 | 0.029 | 0.007 | 0.020 | 0.006 |
| Generation R | 2537 | 0.049 | 0.013 | 0.026 | 0.009 |

SE, standard error. HAPO study is independent of the birth weight meta-analysis whilst NFBC1966 and Generation R (European component) are part of the meta-analysis.

Supplementary Table 10. Look-up of maternal genotype effect on offspring BW (unadjusted for foetal genotype) in up to 68,254 mothers for the 60 BW loci detected in foetal birth weight GWAS.

| Locus | Lead SNP | Allele | | Foetal (European ancestry) GWAS | | Maternal GWAS | | | | | Foetal Beta - Maternal Beta ^a |
|--------------------|-------------|--------|-------|---------------------------------|-------|---------------|---------|-------|----------------------|--------|--|
| | | Effect | Other | β | SE | EAF | β | SE | P-value | N | |
| <i>WNT4-ZBTB40</i> | rs2473248 | C | T | 0.033 | 0.006 | 0.88 | 0.013 | 0.010 | 0.20 | 48,632 | 0.020 |
| <i>ZBTB7B</i> | rs3753639 | C | T | 0.031 | 0.004 | 0.25 | 0.017 | 0.008 | 0.03 | 48,632 | 0.014 |
| <i>FCGR2B</i> | rs72480273 | C | A | 0.031 | 0.005 | 0.19 | 0.013 | 0.008 | 0.11 | 48,632 | 0.018 |
| <i>DTL</i> | rs61830764 | A | G | 0.022 | 0.004 | 0.37 | -0.002 | 0.007 | 0.76 | 48,632 | 0.020 |
| <i>ATAD2B</i> | rs7575873 | A | G | 0.038 | 0.004 | 0.87 | -0.011 | 0.010 | 0.27 | 48,632 | 0.028 |
| <i>EPAS1</i> | rs1374204 | T | C | 0.047 | 0.004 | 0.69 | 0.024 | 0.006 | 5.6x10 ⁻⁵ | 68,252 | 0.023 |
| <i>PTH1R</i> | rs2242116 | A | G | 0.022 | 0.004 | 0.37 | 0.019 | 0.006 | 9.3x10 ⁻⁴ | 67,542 | 0.004 |
| <i>ADCY5</i> | rs11719201 | T | C | 0.046 | 0.004 | 0.25 | -0.001 | 0.008 | 0.85 | 48,632 | 0.045 |
| <i>CPA3</i> | rs10935733 | T | C | 0.022 | 0.004 | 0.40 | 0.015 | 0.007 | 0.02 | 48,632 | 0.007 |
| <i>CCNL1-LEKR1</i> | rs13322435 | A | G | 0.053 | 0.004 | 0.59 | 0.026 | 0.006 | 9.5x10 ⁻⁶ | 68,250 | 0.027 |
| <i>LCORL</i> | rs925098 | G | A | 0.034 | 0.004 | 0.28 | 0.025 | 0.006 | 2.9x10 ⁻⁵ | 68,254 | 0.009 |
| <i>HHIP</i> | rs6537307 | G | A | 0.025 | 0.004 | 0.49 | 0.019 | 0.005 | 4.4x10 ⁻⁴ | 68,247 | 0.006 |
| <i>5q11.2</i> | rs854037 | A | G | 0.027 | 0.005 | 0.80 | 0.009 | 0.007 | 0.18 | 68,219 | 0.018 |
| <i>EBF1</i> | rs7729301 | A | G | 0.024 | 0.004 | 0.74 | 0.039 | 0.007 | 8.9x10 ⁻⁸ | 48,632 | -0.015 |
| <i>CDKAL1</i> | rs35261542 | C | A | 0.044 | 0.004 | 0.74 | 0.004 | 0.007 | 0.60 | 48,632 | 0.040 |
| <i>HIST1H2BE</i> | rs9379832 | A | G | 0.023 | 0.004 | 0.73 | 0.008 | 0.007 | 0.30 | 48,632 | 0.015 |
| <i>HMGA1</i> | rs7742369 | G | A | 0.028 | 0.005 | 0.19 | 0.017 | 0.007 | 0.01 | 68,253 | 0.011 |
| <i>L3MBTL3</i> | rs1415701 | G | A | 0.025 | 0.004 | 0.73 | 0.037 | 0.006 | 3.7x10 ⁻⁹ | 66,897 | -0.012 |
| <i>ESR1</i> | rs1101081 | C | T | 0.038 | 0.004 | 0.72 | 0.021 | 0.007 | 3.8x10 ⁻³ | 48,632 | 0.017 |
| <i>GNA12</i> | rs798489 | C | T | 0.023 | 0.004 | 0.72 | 0.023 | 0.006 | 1.6x10 ⁻⁴ | 68,253 | 0.001 |
| <i>IGF2BP3</i> | rs11765649 | T | C | 0.027 | 0.004 | 0.74 | 0.015 | 0.007 | 0.04 | 48,632 | 0.012 |
| <i>TBX20</i> | rs6959887 | A | G | 0.023 | 0.004 | 0.61 | 0.009 | 0.006 | 0.10 | 68,232 | 0.014 |
| <i>YKT6-GCK</i> | rs138715366 | C | T | 0.241 | 0.023 | 0.99 | 0.137 | 0.038 | 2.8x10 ⁻⁴ | 48,632 | 0.104 |
| <i>MLXIPL</i> | rs62466330 | C | T | 0.051 | 0.011 | 0.07 | 0.022 | 0.013 | 0.08 | 48,632 | 0.029 |
| <i>ANK1-NKX6-3</i> | rs13266210 | A | G | 0.031 | 0.005 | 0.78 | 0.005 | 0.007 | 0.43 | 68,252 | 0.026 |
| <i>TRIB1</i> | rs6989280 | G | A | 0.022 | 0.004 | 0.72 | 0.020 | 0.006 | 1.3x10 ⁻³ | 67,520 | 0.003 |
| <i>SLC45A4</i> | rs12543725 | G | A | 0.023 | 0.004 | 0.58 | 0.021 | 0.005 | 1.4x10 ⁻⁴ | 68,222 | 0.002 |
| <i>PTCH1</i> | rs28510415 | G | A | 0.056 | 0.007 | 0.09 | 0.034 | 0.011 | 2.2x10 ⁻³ | 48,632 | 0.022 |
| <i>LPAR1</i> | rs2150052 | T | A | 0.021 | 0.004 | 0.50 | 0.017 | 0.005 | 1.4x10 ⁻³ | 68,245 | 0.004 |
| <i>PHF19</i> | rs7847628 | G | A | 0.023 | 0.004 | 0.68 | 0.009 | 0.007 | 0.20 | 48,632 | 0.014 |
| <i>STRBP</i> | rs700059 | G | A | 0.033 | 0.005 | 0.16 | 0.018 | 0.008 | 0.02 | 68,251 | 0.016 |

| Locus | Lead SNP | Allele | | Foetal (European ancestry) GWAS | | Maternal GWAS | | | | | Foetal Beta - Maternal Beta ^a |
|----------------------|-------------|--------|-------|---------------------------------|-------|---------------|---------|-------|-----------------------|--------|--|
| | | Effect | Other | β | SE | EAF | β | SE | P-value | N | |
| <i>HHEX-IDE</i> | rs61862780 | T | C | 0.028 | 0.004 | 0.49 | -0.004 | 0.006 | 0.55 | 48,632 | 0.024 |
| <i>NT5C2</i> | rs74233809 | C | T | 0.037 | 0.007 | 0.08 | 0.005 | 0.012 | 0.70 | 48,632 | 0.032 |
| <i>ADRB1</i> | rs7076938 | T | C | 0.036 | 0.004 | 0.73 | 0.010 | 0.006 | 0.09 | 68,254 | 0.026 |
| <i>PLEKHA1</i> | rs2421016 | T | C | 0.021 | 0.004 | 0.48 | 0.016 | 0.005 | 3.3×10^{-3} | 68,253 | 0.005 |
| <i>INS-IGF2</i> | rs72851023 | T | C | 0.048 | 0.008 | 0.08 | 0.015 | 0.012 | 0.24 | 48,632 | 0.033 |
| <i>MTNR1B</i> | rs10830963 | G | C | 0.023 | 0.004 | 0.29 | 0.048 | 0.006 | 5.1×10^{-15} | 67,603 | -0.025 |
| <i>APOLD1</i> | rs11055034 | C | A | 0.022 | 0.004 | 0.71 | 0.002 | 0.006 | 0.74 | 67,597 | 0.020 |
| <i>ABCC9</i> | rs139975827 | G | A | 0.027 | 0.007 | 0.63 | -0.010 | 0.007 | 0.13 | 48,632 | 0.017 |
| <i>ITPR2</i> | rs12823128 | T | C | 0.021 | 0.004 | 0.53 | 0.009 | 0.005 | 0.08 | 68,187 | 0.012 |
| <i>HMGA2</i> | rs1351394 | T | C | 0.044 | 0.004 | 0.49 | 0.034 | 0.005 | 1.4×10^{-10} | 68,247 | 0.010 |
| <i>IGF1</i> | rs7964361 | A | G | 0.039 | 0.007 | 0.10 | 0.018 | 0.010 | 0.06 | 68,251 | 0.021 |
| <i>LINC00332</i> | rs2324499 | G | C | 0.022 | 0.004 | 0.68 | 0.012 | 0.006 | 0.05 | 68,213 | 0.011 |
| <i>RB1</i> | rs2854355 | G | A | 0.023 | 0.004 | 0.26 | 0.017 | 0.007 | 0.03 | 48,632 | 0.007 |
| <i>RNF219-AS1</i> | rs1819436 | C | T | 0.033 | 0.006 | 0.88 | 0.007 | 0.010 | 0.50 | 48,632 | 0.026 |
| <i>FES</i> | rs12906125 | G | A | 0.023 | 0.004 | 0.68 | 0.029 | 0.007 | 1.8×10^{-5} | 48,632 | -0.006 |
| <i>IGF1R</i> | rs7402982 | A | G | 0.023 | 0.004 | 0.41 | 0.021 | 0.007 | 1.7×10^{-3} | 48,632 | 0.002 |
| <i>GPR139</i> | rs1011939 | G | A | 0.022 | 0.004 | 0.28 | 0.007 | 0.007 | 0.34 | 48,632 | 0.015 |
| <i>CLDN7</i> | rs113086489 | T | C | 0.031 | 0.004 | 0.54 | 0.022 | 0.007 | 7.7×10^{-4} | 48,632 | 0.009 |
| <i>SUZ12P1-CRLF3</i> | rs144843919 | G | A | 0.066 | 0.012 | 0.96 | 0.041 | 0.018 | 0.02 | 48,632 | 0.025 |
| <i>SP6-SP2</i> | rs12942207 | C | T | 0.022 | 0.004 | 0.30 | 0.010 | 0.007 | 0.14 | 48,632 | 0.012 |
| <i>ACTL9</i> | rs61154119 | T | G | 0.028 | 0.005 | 0.85 | 0.047 | 0.009 | 1.5×10^{-7} | 48,632 | -0.019 |
| <i>PEPD</i> | rs10402712 | A | G | 0.022 | 0.004 | 0.26 | -0.002 | 0.007 | 0.76 | 48,632 | 0.020 |
| <i>JAG1</i> | rs6040076 | C | G | 0.023 | 0.004 | 0.51 | 0.015 | 0.007 | 0.02 | 48,632 | 0.008 |
| <i>C20orf203</i> | rs28530618 | A | G | 0.026 | 0.004 | 0.47 | 0.003 | 0.006 | 0.62 | 48,632 | 0.023 |
| <i>MAFB</i> | rs6016377 | T | C | 0.024 | 0.004 | 0.45 | 0.005 | 0.006 | 0.35 | 67,909 | 0.019 |
| <i>NRIP1</i> | rs2229742 | G | C | 0.036 | 0.006 | 0.88 | 0.018 | 0.009 | 0.04 | 68,155 | 0.018 |
| <i>KREMEN1</i> | rs134594 | C | T | 0.023 | 0.004 | 0.36 | 0.004 | 0.006 | 0.53 | 66,634 | 0.020 |
| <i>SREBF2</i> | rs62240962 | C | T | 0.047 | 0.007 | 0.92 | 0.027 | 0.012 | 0.02 | 48,632 | 0.020 |
| <i>PLAC1</i> | rs11096402 | G | A | 0.028 | 0.005 | 0.24 | 0.001 | 0.003 | 0.82 | 45,353 | 0.027 |

SE, standard error, EAF, effect allele frequency. The lead SNPs at *MTNR1B*, *HMGA2* and *L3MBTL3* attained genome-wide significance in the maternal, as well as the foetal GWAS. ^aFoetal GWAS effect size subtracted from maternal GWAS effect size: 55/60 beta values were more positive in the foetal GWAS (aligned to BW-raising allele) than in the maternal GWAS (2-tailed binomial sign test $P=1 \times 10^{-11}$).

Supplementary Table 11. Maternal and foetal conditional analyses in 12,909 mother-child pairs.

| Locus | Index variant | EA/ NEA | Maternal effect on offspring birth weight | | | | | | Foetal effect on birth weight | | | | | |
|--------------------|---------------|------------|---|-------|----------------------|------------------------------|-------|----------------------|-------------------------------|-------|-----------------------|--------------------------------|-------|-----------------------|
| | | | Unadjusted | | | Adjusted for foetal genotype | | | Unadjusted | | | Adjusted for maternal genotype | | |
| | | | β | SE | P-value | β | SE | P-value | β | SE | P-value | β | SE | P-value |
| <i>WNT4-ZBTB40</i> | rs2473248 | C/T | 0.004 | 0.018 | 0.82 | -0.006 | 0.020 | 0.76 | 0.033 | 0.018 | 0.06 | 0.039 | 0.020 | 0.05 |
| <i>ZBTB7B</i> | rs3753639 | C/T | 0.022 | 0.015 | 0.16 | -0.005 | 0.017 | 0.79 | 0.060 | 0.016 | 9.8×10^{-5} | 0.060 | 0.018 | 5.7×10^{-4} |
| <i>FCGR2B</i> | rs72480273 | C/A | 0.034 | 0.017 | 0.05 | 0.018 | 0.019 | 0.34 | 0.042 | 0.017 | 0.01 | 0.031 | 0.019 | 0.11 |
| <i>DTL</i> | rs61830764 | A/G | 0.026 | 0.014 | 0.06 | 0.016 | 0.016 | 0.31 | 0.029 | 0.014 | 0.04 | 0.027 | 0.016 | 0.09 |
| <i>ATAD2B</i> | rs7575873 | A/G | 0.018 | 0.019 | 0.35 | 0.007 | 0.022 | 0.74 | 0.042 | 0.019 | 0.03 | 0.047 | 0.022 | 0.03 |
| <i>EPAS1</i> | rs1374204 | T/C | 0.032 | 0.013 | 0.02 | 0.008 | 0.015 | 0.60 | 0.056 | 0.013 | 2.6×10^{-5} | 0.047 | 0.015 | 1.9×10^{-3} |
| <i>PTH1R</i> | rs2242116 | A/G | 0.025 | 0.013 | 0.05 | 0.031 | 0.015 | 0.03 | 0.002 | 0.013 | 0.86 | -0.013 | 0.014 | 0.38 |
| <i>ADCY5</i> | rs11719201 | T/C | 0.000 | 0.015 | 1.00 | -0.025 | 0.017 | 0.15 | 0.057 | 0.015 | 2.2×10^{-4} | 0.075 | 0.017 | 1.5×10^{-5} |
| <i>CPA3</i> | rs10935733 | T/C | -0.004 | 0.013 | 0.73 | -0.006 | 0.014 | 0.66 | 0.007 | 0.013 | 0.57 | 0.013 | 0.014 | 0.36 |
| <i>CCNL1-LEKR1</i> | rs13322435 | A/G | 0.028 | 0.013 | 0.03 | -0.015 | 0.014 | 0.31 | 0.091 | 0.013 | 7.9×10^{-13} | 0.097 | 0.014 | 9.7×10^{-12} |
| <i>LCORL</i> | rs925098 | G/A | 0.012 | 0.014 | 0.39 | -0.017 | 0.016 | 0.29 | 0.057 | 0.014 | 3.1×10^{-5} | 0.060 | 0.015 | 1.0×10^{-4} |
| <i>HHIP</i> | rs6537307 | G/A | 0.028 | 0.013 | 0.03 | 0.003 | 0.014 | 0.85 | 0.057 | 0.013 | 1.1×10^{-5} | 0.057 | 0.015 | 7.8×10^{-5} |
| <i>5q11.2</i> | rs854037 | A/G | 0.019 | 0.015 | 0.20 | 0.016 | 0.017 | 0.35 | 0.022 | 0.015 | 0.15 | 0.011 | 0.017 | 0.52 |
| <i>EBF1</i> | rs7729301 | A/G | 0.036 | 0.014 | 8.8×10^{-3} | 0.040 | 0.015 | 9.6×10^{-3} | 0.024 | 0.014 | 0.08 | 0.010 | 0.016 | 0.52 |
| <i>CDKAL1</i> | rs35261542 | C/A | 0.003 | 0.014 | 0.81 | -0.028 | 0.016 | 0.07 | 0.057 | 0.014 | 4.3×10^{-5} | 0.071 | 0.016 | 6.1×10^{-6} |
| <i>HIST1H2BE</i> | rs9379832 | A/G | -0.005 | 0.014 | 0.74 | -0.012 | 0.016 | 0.45 | 0.022 | 0.014 | 0.12 | 0.026 | 0.016 | 0.10 |
| <i>HMGA1</i> | rs7742369 | G/A | 0.027 | 0.015 | 0.08 | 0.006 | 0.017 | 0.72 | 0.043 | 0.015 | 4.9×10^{-3} | 0.035 | 0.017 | 0.04 |
| <i>L3MBTL3</i> | rs1415701 | G/A | 0.035 | 0.014 | 0.01 | 0.019 | 0.016 | 0.22 | 0.052 | 0.014 | 2.7×10^{-4} | 0.041 | 0.016 | 0.01 |
| <i>ESR1</i> | rs1101081 | C/T | 0.009 | 0.014 | 0.51 | -0.004 | 0.016 | 0.82 | 0.029 | 0.014 | 0.04 | 0.033 | 0.016 | 0.04 |
| <i>GNA12</i> | rs798489 | C/T | 0.018 | 0.015 | 0.22 | 0.011 | 0.017 | 0.50 | 0.034 | 0.015 | 0.02 | 0.029 | 0.017 | 0.08 |
| <i>IGF2BP3</i> | rs11765649 | T/C | 0.007 | 0.015 | 0.64 | 0.006 | 0.017 | 0.74 | 0.017 | 0.015 | 0.25 | 0.013 | 0.017 | 0.45 |
| <i>TBX20</i> | rs6959887 | A/G | 0.018 | 0.013 | 0.15 | -0.002 | 0.014 | 0.89 | 0.040 | 0.013 | 1.7×10^{-3} | 0.036 | 0.014 | 0.01 |
| <i>YKT6-GCK</i> | rs138715366 | C/T | -0.030 | 0.087 | 0.73 | -0.147 | 0.097 | 0.13 | 0.186 | 0.087 | 0.03 | 0.230 | 0.098 | 0.02 |
| <i>MLXIPL</i> | rs62466330 | C/T | 0.019 | 0.028 | 0.50 | -0.015 | 0.031 | 0.62 | 0.104 | 0.028 | 2.2×10^{-4} | 0.129 | 0.031 | 3.3×10^{-5} |
| <i>ANK1-NKX6-3</i> | rs13266210 | A/G | 0.013 | 0.015 | 0.41 | 0.009 | 0.017 | 0.61 | 0.018 | 0.015 | 0.24 | 0.023 | 0.017 | 0.18 |
| <i>TRIB1</i> | rs6989280 | G/A | 0.014 | 0.014 | 0.32 | 0.019 | 0.016 | 0.24 | -0.003 | 0.014 | 0.84 | -0.012 | 0.016 | 0.46 |
| <i>SLC45A4</i> | rs12543725 | G/A | 0.012 | 0.013 | 0.36 | 0.004 | 0.015 | 0.79 | 0.013 | 0.013 | 0.31 | 0.014 | 0.015 | 0.34 |
| <i>PTCH1</i> | rs28510415 | G/A | 0.037 | 0.023 | 0.11 | -0.005 | 0.026 | 0.84 | 0.105 | 0.023 | 8.0×10^{-6} | 0.105 | 0.027 | 7.2×10^{-5} |
| <i>LPAR1</i> | rs2150052 | T/A | 0.019 | 0.013 | 0.13 | 0.008 | 0.014 | 0.56 | 0.017 | 0.013 | 0.17 | 0.010 | 0.014 | 0.47 |
| <i>PHF19</i> | rs7847628 | G/A | 0.020 | 0.013 | 0.13 | 0.018 | 0.015 | 0.24 | 0.022 | 0.013 | 0.11 | 0.017 | 0.015 | 0.25 |
| <i>STRBP</i> | rs700059 | G/A | 0.017 | 0.016 | 0.27 | -0.006 | 0.018 | 0.76 | 0.053 | 0.016 | 8.0×10^{-4} | 0.062 | 0.018 | 4.5×10^{-4} |
| <i>HHEX-IDE</i> | rs61862780 | T/C | 0.012 | 0.013 | 0.36 | -0.015 | 0.014 | 0.29 | 0.053 | 0.013 | 3.0×10^{-5} | 0.059 | 0.014 | 3.1×10^{-5} |

| Locus | Index variant | EA/ NEA | Maternal effect on offspring birth weight | | | | | | Foetal effect on birth weight | | | | | |
|--------------------------|---------------|------------|---|-------|----------------------|------------------------------|-------|---------|-------------------------------|-------|----------------------|--------------------------------|-------|----------------------|
| | | | Unadjusted | | | Adjusted for foetal genotype | | | Unadjusted | | | Adjusted for maternal genotype | | |
| | | | β | SE | P-value | β | SE | P-value | β | SE | P-value | β | SE | P-value |
| <i>NT5C2</i> | rs74233809 | C/T | -0.002 | 0.021 | 0.94 | -0.010 | 0.024 | 0.67 | 0.016 | 0.021 | 0.44 | 0.011 | 0.024 | 0.64 |
| <i>ADRB1</i> | rs7076938 | T/C | 0.045 | 0.014 | 8.8×10^{-4} | 0.032 | 0.015 | 0.03 | 0.050 | 0.014 | 2.6×10^{-4} | 0.037 | 0.015 | 0.02 |
| <i>PLEKHA1</i> | rs2421016 | T/C | 0.017 | 0.012 | 0.17 | 0.018 | 0.014 | 0.19 | 0.006 | 0.012 | 0.63 | 0.001 | 0.014 | 0.95 |
| <i>INS-IGF2</i> | rs72851023 | T/C | 0.031 | 0.027 | 0.26 | 0.008 | 0.031 | 0.79 | 0.060 | 0.028 | 0.03 | 0.061 | 0.031 | 0.05 |
| <i>MTNR1B</i> | rs10830963 | G/C | 0.047 | 0.015 | 1.7×10^{-3} | 0.035 | 0.017 | 0.04 | 0.049 | 0.015 | 9.8×10^{-4} | 0.032 | 0.017 | 0.06 |
| <i>APOLD1</i> | rs11055034 | C/A | -0.003 | 0.014 | 0.83 | -0.013 | 0.016 | 0.42 | 0.005 | 0.014 | 0.72 | 0.011 | 0.016 | 0.51 |
| <i>ABCC9</i> | rs139975827 | G/A | 0.005 | 0.014 | 0.72 | -0.011 | 0.015 | 0.47 | 0.033 | 0.014 | 0.02 | 0.041 | 0.015 | 0.01 |
| <i>ITPR2</i> | rs12823128 | T/C | -0.003 | 0.013 | 0.83 | -0.016 | 0.015 | 0.28 | 0.022 | 0.013 | 0.10 | 0.024 | 0.015 | 0.10 |
| <i>HMGA2</i> | rs1351394 | T/C | 0.023 | 0.013 | 0.07 | 0.007 | 0.015 | 0.65 | 0.038 | 0.013 | 3.0×10^{-3} | 0.038 | 0.014 | 0.01 |
| <i>IGF1</i> | rs7964361 | A/G | -0.005 | 0.023 | 0.85 | -0.016 | 0.026 | 0.55 | 0.025 | 0.023 | 0.28 | 0.030 | 0.026 | 0.25 |
| <i>LINC00332</i> | rs2324499 | G/C | 0.006 | 0.013 | 0.68 | 0.001 | 0.015 | 0.96 | 0.019 | 0.013 | 0.15 | 0.019 | 0.015 | 0.22 |
| <i>RB1</i> | rs2854355 | G/A | -0.013 | 0.015 | 0.37 | -0.027 | 0.016 | 0.10 | 0.020 | 0.015 | 0.18 | 0.029 | 0.016 | 0.07 |
| <i>RNF219-AS1</i> | rs1819436 | C/T | 0.040 | 0.018 | 0.03 | 0.013 | 0.020 | 0.52 | 0.057 | 0.018 | 1.8×10^{-3} | 0.046 | 0.020 | 0.02 |
| <i>FES</i> | rs12906125 | G/A | 0.006 | 0.014 | 0.66 | -0.004 | 0.016 | 0.82 | 0.016 | 0.014 | 0.27 | 0.012 | 0.016 | 0.47 |
| <i>IGF1R</i> | rs7402982 | A/G | 0.015 | 0.013 | 0.24 | 0.003 | 0.015 | 0.83 | 0.025 | 0.013 | 0.06 | 0.027 | 0.015 | 0.07 |
| <i>GPR139</i> | rs1011939 | G/A | 0.015 | 0.014 | 0.27 | 0.013 | 0.015 | 0.39 | 0.008 | 0.014 | 0.57 | 0.000 | 0.015 | 1.00 |
| <i>CLDN7</i> | rs113086489 | T/C | 0.018 | 0.013 | 0.16 | 0.002 | 0.014 | 0.89 | 0.043 | 0.013 | 5.7×10^{-4} | 0.044 | 0.014 | 1.9×10^{-3} |
| <i>SUZ12P1-CRLF3</i> | rs144843919 | G/A | 0.003 | 0.037 | 0.93 | -0.030 | 0.041 | 0.47 | 0.045 | 0.038 | 0.24 | 0.047 | 0.043 | 0.27 |
| <i>SP6-SP2</i> | rs12942207 | C/T | -0.001 | 0.014 | 0.92 | -0.003 | 0.016 | 0.82 | 0.007 | 0.014 | 0.63 | 0.008 | 0.015 | 0.59 |
| <i>ACTL9</i> | rs61154119 | T/G | -0.005 | 0.016 | 0.75 | -0.022 | 0.018 | 0.23 | 0.032 | 0.016 | 0.05 | 0.041 | 0.018 | 0.02 |
| <i>PEPD</i> | rs10402712 | A/G | -0.013 | 0.014 | 0.36 | -0.028 | 0.016 | 0.07 | 0.016 | 0.014 | 0.26 | 0.025 | 0.016 | 0.11 |
| <i>JAG1</i> | rs6040076 | C/G | 0.000 | 0.013 | 0.98 | -0.007 | 0.015 | 0.64 | 0.010 | 0.013 | 0.43 | 0.017 | 0.015 | 0.24 |
| <i>C20orf203</i> | rs28530618 | A/G | 0.015 | 0.013 | 0.23 | 0.013 | 0.014 | 0.34 | 0.006 | 0.013 | 0.63 | 0.003 | 0.014 | 0.85 |
| <i>MAFB</i> | rs6016377 | T/C | 0.000 | 0.013 | 0.99 | -0.021 | 0.015 | 0.16 | 0.042 | 0.013 | 1.4×10^{-3} | 0.058 | 0.015 | 9.9×10^{-5} |
| <i>NRIP1</i> | rs2229742 | G/C | 0.026 | 0.023 | 0.26 | 0.018 | 0.026 | 0.49 | 0.031 | 0.023 | 0.17 | 0.018 | 0.026 | 0.49 |
| <i>KREMEN1</i> | rs134594 | C/T | 0.016 | 0.013 | 0.22 | 0.007 | 0.015 | 0.64 | 0.017 | 0.013 | 0.20 | 0.015 | 0.015 | 0.30 |
| <i>SREBF2</i> | rs62240962 | C/T | 0.019 | 0.026 | 0.47 | 0.005 | 0.029 | 0.87 | 0.049 | 0.026 | 0.06 | 0.049 | 0.029 | 0.09 |
| <i>PLAC1^a</i> | rs11096402 | G/A | 0.009 | 0.022 | 0.68 | -0.004 | 0.024 | 0.88 | 0.013 | 0.024 | 0.60 | 0.014 | 0.027 | 0.61 |

^aFor analysis of the *PLAC1* locus on the X chromosome, we used female offsprings only: N = 4,940 mother-child pairs

EA: Effect Allele. NEA: Non Effect Allele. SE: standard error. Effects (beta values) are aligned to the BW-raising allele as reported in Extended Data Table 1a.

Supplementary Table 12. Summary results of LD Score regression analyses between birth weight and various diseases, metabolic and anthropometric traits.

(a) Birth weight and other traits

| Phenotype1 | Phenotype2 | Genetic Correlation (r_g) | SE | z | P-value | Sample size of phenotype 2 | Number of SNPs | Reference (PMID) |
|--------------|---|-------------------------------|-------|-------|-----------------------|----------------------------|----------------|------------------|
| Birth weight | Infant head circumference | 0.389 | 0.073 | 5.30 | 1.2×10^{-7} | 10,770 | 1,019,674 | 22504419 |
| Birth weight | Birth length | 0.805 | 0.058 | 13.98 | 2.0×10^{-44} | 28,460 | 983,823 | 25281659 |
| Birth weight | Height (2014) | 0.406 | 0.027 | 15.18 | 4.8×10^{-52} | 253,300 | 1,049,262 | 25282103 |
| Birth weight | Height (2010) | 0.401 | 0.028 | 14.09 | 4.6×10^{-45} | 133,900 | 1,026,905 | 20881960 |
| Birth weight | Height (women) | 0.400 | 0.032 | 12.59 | 2.4×10^{-36} | 73,140 | 1,028,514 | 23754948 |
| Birth weight | Height (men) | 0.384 | 0.035 | 11.00 | 3.7×10^{-28} | 60,590 | 1,024,139 | 23754948 |
| Birth weight | Extreme height | 0.399 | 0.043 | 9.20 | 3.5×10^{-20} | 16,200 | 739,427 | 23563607 |
| Birth weight | Body mass index (2015) | 0.114 | 0.025 | 4.49 | 7.3×10^{-6} | 322,200 | 1,050,986 | 25673413 |
| Birth weight | Body mass index (2010) | 0.120 | 0.033 | 3.66 | 3.0×10^{-4} | 123,900 | 1,028,235 | 20935630 |
| Birth weight | Extreme Body mass index | 0.116 | 0.048 | 2.40 | 0.017 | 16,070 | 738,850 | 23563607 |
| Birth weight | Body mass index (men) | 0.148 | 0.040 | 3.67 | 2.4×10^{-4} | 58,670 | 1,025,063 | 23754948 |
| Birth weight | Body mass index (women) | 0.096 | 0.035 | 2.76 | 0.006 | 67,960 | 1,029,327 | 23754948 |
| Birth weight | Weight (men) | 0.377 | 0.039 | 9.57 | 1.1×10^{-21} | 58,350 | 1,025,155 | 23754948 |
| Birth weight | Weight (women) | 0.280 | 0.036 | 7.70 | 1.4×10^{-14} | 67,590 | 1,029,320 | 23754948 |
| Birth weight | Waist-hip ratio (2015) | -0.052 | 0.031 | -1.66 | 0.097 | 212,200 | 1,048,211 | 25673412 |
| Birth weight | Waist-hip ratio (adjusted for BMI - 2015) | -0.159 | 0.032 | -4.93 | 8.3×10^{-7} | 210,100 | 1,051,567 | 25673412 |
| Birth weight | Waist-hip ratio (adjusted for BMI - 2010) | -0.189 | 0.041 | -4.59 | 4.4×10^{-6} | 77,220 | 1,027,370 | 20935629 |
| Birth weight | Waist-hip ratio (men) | -0.080 | 0.053 | -1.51 | 0.132 | 34,640 | 1,025,922 | 23754948 |
| Birth weight | Waist-hip ratio (women) | -0.060 | 0.054 | -1.11 | 0.268 | 42,730 | 1,029,620 | 23754948 |
| Birth weight | Extreme waist-hip ratio | -0.254 | 0.071 | -3.57 | 3.6×10^{-4} | 10,260 | 674,047 | 23563607 |
| Birth weight | Hip circumference | 0.281 | 0.030 | 9.30 | 1.4×10^{-20} | 213,000 | 1,048,111 | 25673412 |
| Birth weight | Hip circumference (adjusted for BMI) | 0.314 | 0.033 | 9.49 | 2.3×10^{-21} | 211,100 | 1,049,874 | 25673412 |
| Birth weight | Hip circumference (men) | 0.339 | 0.063 | 5.34 | 9.3×10^{-8} | 32,850 | 1,030,276 | 23754948 |
| Birth weight | Hip circumference (women) | 0.260 | 0.057 | 4.54 | 5.6×10^{-6} | 40,360 | 1,031,503 | 23754948 |
| Birth weight | Waist circumference | 0.177 | 0.028 | 6.29 | 3.9×10^{-10} | 232,100 | 1,048,634 | 25673412 |
| Birth weight | Waist circumference (adjusted for BMI) | 0.128 | 0.032 | 4.03 | 5.7×10^{-5} | 231,400 | 1,049,871 | 25673412 |
| Birth weight | Waist circumference (men) | 0.218 | 0.053 | 4.11 | 3.9×10^{-5} | 38,310 | 1,025,680 | 23754948 |
| Birth weight | Waist circumference (women) | 0.135 | 0.047 | 2.86 | 0.004 | 47,320 | 1,027,090 | 23754948 |
| Birth weight | Obesity (class 1) | 0.102 | 0.035 | 2.97 | 0.003 | 98,700 | 965,717 | 23563607 |
| Birth weight | Obesity (class 2) | 0.099 | 0.040 | 2.46 | 0.014 | 72,550 | 890,085 | 23563607 |

| Phenotype1 | Phenotype2 | Genetic Correlation (r_g) | SE | z | P-value | Sample size of phenotype 2 | Number of SNPs | Reference (PMID) |
|--------------|--|-------------------------------|-------|-------|-----------------------|----------------------------|----------------|------------------|
| Birth weight | Obesity (class 3) | 0.136 | 0.055 | 2.48 | 0.013 | 50,360 | 621,425 | 23563607 |
| Birth weight | Childhood obesity | 0.192 | 0.049 | 3.95 | 7.8×10^{-5} | 13,850 | 1,057,303 | 22484627 |
| Birth weight | Overweight | 0.061 | 0.033 | 1.83 | 0.067 | 158,900 | 1,012,975 | 23563607 |
| Birth weight | Anorexia nervosa | 0.029 | 0.033 | 0.91 | 0.365 | 17,770 | 1,042,946 | 24514567 |
| Birth weight | Crohn's disease | 0.047 | 0.044 | 1.07 | 0.284 | 20,883 | 1,066,772 | 26192919 |
| Birth weight | Irritable bowel syndrome | 0.009 | 0.052 | 0.18 | 0.858 | 27,432 | 1,091,971 | 26192919 |
| Birth weight | Rheumatoid arthritis | -0.004 | 0.046 | -0.08 | 0.935 | 25,710 | 986,060 | 20453842 |
| Birth weight | Lumbar spine bone mineral density | -0.027 | 0.043 | -0.63 | 0.531 | 31,800 | 1,061,087 | 22504420 |
| Birth weight | Lumbar spine bone mineral density (men) | -0.098 | 0.064 | -1.53 | 0.127 | 9,980 | 1,060,919 | 22504420 |
| Birth weight | Lumbar spine bone mineral density (women) | -0.003 | 0.044 | -0.07 | 0.943 | 22,180 | 1,064,756 | 22504420 |
| Birth weight | Femoral neck bone mineral density | 0.013 | 0.039 | 0.34 | 0.736 | 32,960 | 1,065,335 | 22504420 |
| Birth weight | Femoral neck bone mineral density (men) | -0.019 | 0.057 | -0.33 | 0.745 | 9,971 | 1,056,535 | 22504420 |
| Birth weight | Femoral neck bone mineral density (women) | 0.020 | 0.050 | 0.40 | 0.686 | 22,990 | 1,069,031 | 22504420 |
| Birth weight | Coronary artery disease | -0.295 | 0.051 | -5.80 | 6.5×10^{-9} | 84,270 | 932,642 | 21378990 |
| Birth weight | Systolic blood pressure (adjusted for BMI) | -0.263 | 0.039 | -6.72 | 1.8×10^{-11} | 69,395 | 990,423 | 21909115 |
| Birth weight | Systolic blood pressure (unadjusted for BMI) | -0.215 | 0.030 | -7.21 | 5.5×10^{-13} | 127,969 | 1,186,583 | UK Biobank data |
| Birth weight | Diastolic blood pressure (adjusted for BMI) | -0.224 | 0.042 | -5.35 | 8.7×10^{-8} | 69,395 | 990,446 | 21909115 |
| Birth weight | Triglycerides (2010) | -0.167 | 0.038 | -4.41 | 1.0×10^{-5} | 96,600 | 1,024,575 | 20686565 |
| Birth weight | Triglycerides (2013) | -0.112 | 0.031 | -3.61 | 3.0×10^{-4} | 177,900 | 1,026,718 | 24097068 |
| Birth weight | Total cholesterol (2010) | -0.127 | 0.040 | -3.21 | 0.001 | 100,200 | 1,026,024 | 20686565 |
| Birth weight | Total cholesterol (2013) | -0.105 | 0.033 | -3.17 | 0.002 | 187,400 | 1,027,605 | 24097068 |
| Birth weight | Low density lipoprotein (2010) | -0.136 | 0.047 | -2.90 | 0.004 | 95,450 | 1,024,730 | 20686565 |
| Birth weight | Low density lipoprotein (2013) | -0.102 | 0.036 | -2.87 | 0.004 | 173,100 | 1,026,705 | 24097068 |
| Birth weight | High density lipoprotein (2010) | 0.087 | 0.039 | 2.23 | 0.026 | 99,900 | 1,026,029 | 20686565 |
| Birth weight | High density lipoprotein (2013) | 0.056 | 0.029 | 1.97 | 0.048 | 187,200 | 1,027,731 | 24097068 |
| Birth weight | Type 2 diabetes | -0.271 | 0.056 | -4.88 | 1.1×10^{-6} | 69,030 | 976,437 | 22885922 |
| Birth weight | Fasting insulin | -0.150 | 0.071 | -2.10 | 0.036 | 46,190 | 1,054,692 | 20081858 |
| Birth weight | Fasting insulin (adjusting for BMI) | -0.178 | 0.055 | -3.26 | 0.001 | 51,750 | 1,121,151 | 22581228 |
| Birth weight | HbA1c | -0.166 | 0.060 | -2.75 | 0.006 | 46,368 | 1,109,233 | 20858683 |
| Birth weight | 2hr glucose | -0.245 | 0.094 | -2.62 | 0.009 | 15,230 | 1,028,518 | 20081857 |
| Birth weight | Fasting glucose | -0.073 | 0.054 | -1.35 | 0.178 | 46,190 | 1,056,257 | 20081858 |
| Birth weight | Fasting glucose (adjusting for BMI) | -0.126 | 0.048 | -2.61 | 0.009 | 58,074 | 1,121,150 | 22581228 |
| Birth weight | HOMA-IR | -0.133 | 0.072 | -1.84 | 0.066 | 46,190 | 1,053,709 | 20081858 |
| Birth weight | HOMA-B | -0.065 | 0.065 | -1.00 | 0.317 | 46,190 | 1,053,581 | 20081858 |

| Phenotype1 | Phenotype2 | Genetic Correlation (r_g) | SE | z | P-value | Sample size of phenotype 2 | Number of SNPs | Reference (PMID) |
|--------------|---|-------------------------------|-------|-------|----------------------|----------------------------|----------------|------------------|
| Birth weight | Educational attainment (college) | 0.111 | 0.042 | 2.63 | 0.009 | 126,600 | 1,017,448 | 25201988 |
| Birth weight | Educational attainment (years) | 0.105 | 0.043 | 2.43 | 0.015 | 126,600 | 1,014,207 | 25201988 |
| Birth weight | Childhood intelligence | 0.123 | 0.075 | 1.64 | 0.100 | 17,990 | 800,741 | 25201988 |
| Birth weight | Autism | -0.089 | 0.053 | -1.69 | 0.091 | 10,263 | 944,654 | 23453885 |
| Birth weight | ADHD | -0.107 | 0.103 | -1.04 | 0.300 | 3,351 | 1,065,753 | 20732625 |
| Birth weight | Major depressive disorder | -0.076 | 0.074 | -1.02 | 0.306 | 18,760 | 887,434 | 22472876 |
| Birth weight | Bipolar disorder | 0.026 | 0.047 | 0.56 | 0.578 | 16,730 | 800,584 | 21926972 |
| Birth weight | Alzheimer's disease | -0.039 | 0.080 | -0.49 | 0.622 | 74,050 | 1,142,332 | 24162737 |
| Birth weight | Schizophrenia | -0.022 | 0.044 | -0.49 | 0.624 | 21,860 | 846,564 | 21926974 |
| Birth weight | Pubertal growth - height at age 10 in females and 12 in males - targets take-off phase of growth spurt | 0.271 | 0.051 | 5.28 | 1.3×10^{-7} | 13,960 | 1,009,067 | 23449627 |
| Birth weight | Pubertal growth - height at age 12 in males | 0.305 | 0.072 | 4.23 | 2.3×10^{-5} | 6,986 | 1,017,307 | 23449627 |
| Birth weight | Pubertal growth - height at age 10 in females | 0.221 | 0.061 | 3.63 | 3.0×10^{-4} | 6,974 | 1,036,441 | 23449627 |
| Birth weight | Pubertal growth - total amount of growth across the pubertal growth period | 0.162 | 0.062 | 2.61 | 0.009 | 10,800 | 1,037,500 | 23449627 |
| Birth weight | Pubertal growth - total amount of growth across the pubertal growth period (males) | 0.173 | 0.080 | 2.15 | 0.032 | 5,043 | 1,039,045 | 23449627 |
| Birth weight | Pubertal growth - total amount of growth across the pubertal growth period (females) | 0.142 | 0.083 | 1.70 | 0.089 | 5,756 | 1,038,734 | 23449627 |
| Birth weight | Pubertal growth - total amount of growth in late adolescence, targeting the timing of peak height growth velocity | 0.109 | 0.075 | 1.46 | 0.145 | 4,282 | 1,039,103 | 23449627 |
| Birth weight | Pubertal growth - total amount of growth in late adolescence (males) | 0.041 | 0.081 | 0.51 | 0.613 | 4,282 | 1,039,103 | 23449627 |
| Birth weight | Pubertal growth - total amount of growth in late adolescence (females) | 0.230 | 0.112 | 2.06 | 0.039 | 4,946 | 1,038,829 | 23449627 |
| Birth weight | Age at menarche (2014) | 0.029 | 0.031 | 0.93 | 0.354 | 133,000 | 1,057,631 | 25231870 |
| Birth weight | Pubertal growth - Tanner scale | -0.137 | 0.125 | -1.10 | 0.273 | 9,918 | 969,224 | 23449627 |
| Birth weight | Pubertal growth - Tanner scale (males) | -0.453 | 0.650 | -0.97 | 0.486 | 3,769 | 969,224 | 23449627 |
| Birth weight | Pubertal growth - Tanner scale (females) | -0.037 | 0.138 | -0.27 | 0.786 | 6,149 | 969,224 | 23449627 |
| Birth weight | Asthma | 0.050 | 0.069 | 0.72 | 0.471 | 10,770 | 1,019,674 | 20860503 |
| Birth weight | Ever smoked | 0.007 | 0.045 | 0.15 | 0.878 | 28,460 | 983,823 | 20418890 |
| Birth weight | Former smoker | 0.097 | 0.061 | 1.58 | 0.114 | 253,300 | 1,049,262 | 20418890 |
| Birth weight | Cigarettes per day | -0.042 | 0.064 | -0.67 | 0.505 | 133,900 | 1,026,905 | 20418890 |
| Birth weight | Smoking start age | 0.041 | 0.087 | 0.47 | 0.636 | 73,140 | 1,028,514 | 20418890 |

(b) Glucose-Related Phenotypes

| Phenotype1 | Phenotype2 | Genetic Correlation (rg) | SE | z | P-value | Sample size of phenotype 2 | Number of SNPs | Reference (PMID) |
|-----------------|-----------------|--------------------------|-------|-------|----------------------|----------------------------|----------------|------------------|
| Type 2 diabetes | Fasting glucose | 0.587 | 0.097 | 6.07 | 1.3×10^{-9} | 46,190 | 1,056,257 | 20081858 |
| Type 2 diabetes | 2hr glucose | 0.301 | 0.12 | 2.47 | 0.014 | 15,230 | 1,028,518 | 20081857 |
| Type 2 diabetes | Fasting insulin | 0.435 | 0.12 | 3.56 | 4.0×10^{-4} | 46,190 | 1,054,692 | 20081858 |
| Type 2 diabetes | HOMA-B | -0.002 | 0.11 | -0.01 | 0.989 | 46,190 | 1,053,581 | 20081858 |

SE, standard error; BMI, body mass index; ADHD, attention deficit hyperactive disorder; HOMA-IR, homeostasis model assessment of insulin resistance; HOMA-B, homeostasis model assessment of beta cell function.

Supplementary Table 13. Estimating the proportion of the BW-adult phenotype covariance attributable to genotyped SNPs in UK Biobank and the Northern Finland Birth Cohort (NFBC) 1966 data.

| Adult phenotype | N ^a | Phenotypic correlation with BW | Genetic correlation with BW (SE) | Residual correlation with BW (SE) | Genetic covariance with BW, gcov (SE) | Residual covariance with BW, rcov (SE) | Proportion of covariance explained by genetic variants directly genotyped: gcov/(gcov+rcov) ^b (95% CI) |
|--|----------------|--------------------------------|----------------------------------|-----------------------------------|---------------------------------------|--|---|
| UK Biobank | | | | | | | |
| Type 2 diabetes | 56,654 | -0.034 | -0.124 (0.081) | -0.001 (0.010) | -0.012 (0.008) | -0.001 (0.008) | 0.96 (0.36, 1.56) |
| Systolic blood pressure (unadjusted for BMI) | 57,581 | -0.072 | -0.201 (0.037) | -0.011 (0.011) | -0.045 (0.008) | -0.008 (0.008) | 0.85 (0.70, 0.99) |
| Systolic blood pressure (adjusted for BMI) | 57,319 | -0.084 | -0.216 (0.030) | -0.042 (0.010) | -0.052 (0.007) | -0.031 (0.007) | 0.62 (0.54, 0.71) |
| Diastolic blood pressure (adjusted for BMI) | 57,230 | -0.072 | -0.156 (0.030) | -0.045 (0.010) | -0.038 (0.007) | -0.033 (0.007) | 0.54 (0.44, 0.64) |
| Coronary artery disease | 57,715 | -0.002 | -0.174 (0.182) | 0.004 (0.009) | -0.008 (0.007) | 0.003 (0.008) | N/A |
| Height | 57,638 | 0.228 | 0.332 (0.026) | 0.194 (0.015) | 0.106 (0.008) | 0.091 (0.007) | 0.54 (0.50, 0.58) |
| Height (men) | 23,385 | 0.202 | 0.343 (0.060) | 0.214 (0.041) | 0.108 (0.019) | 0.090 (0.018) | 0.55 (0.45, 0.64) |
| Height (women) | 34,253 | 0.2 | 0.367 (0.037) | 0.174 (0.029) | 0.131 (0.013) | 0.072 (0.012) | 0.65 (0.58, 0.71) |
| Weight (men) | 23,371 | 0.15 | 0.199 (0.074) | 0.118 (0.027) | 0.050 (0.018) | 0.080 (0.018) | 0.38 (0.24, 0.53) |
| Weight (women) | 34,233 | 0.094 | 0.206 (0.049) | 0.058 (0.019) | 0.056 (0.013) | 0.040 (0.013) | 0.58 (0.45, 0.72) |
| Body mass index | 57,402 | 0.043 | 0.063 (0.036) | 0.031 (0.011) | 0.014 (0.008) | 0.023 (0.008) | 0.39 (0.17, 0.61) |
| Body mass index (men) | 34,077 | 0.045 | 0.049 (0.051) | 0.019 (0.019) | 0.013 (0.013) | 0.013 (0.013) | 0.49 (-0.01, 0.98) |
| Body mass index (women) | 23,325 | 0.024 | 0.072 (0.087) | 0.037 (0.025) | 0.015 (0.018) | 0.028 (0.018) | 0.35 (-0.06, 0.77) |
| Obesity (class 1) | 33,622 | 0.048 | 0.080 (0.051) | 0.040 (0.020) | 0.021 (0.013) | 0.027 (0.013) | 0.44 (0.17, 0.71) |

| Adult phenotype | N ^a | Phenotypic correlation with BW | Genetic correlation with BW (SE) | Residual correlation with BW (SE) | Genetic covariance with BW, gcov (SE) | Residual covariance with BW, rcov (SE) | Proportion of covariance explained by genetic variants directly genotyped: gcov/(gcov+rcov) ^b (95% CI) |
|--|----------------|--------------------------------|----------------------------------|-----------------------------------|---------------------------------------|--|---|
| Obesity (class 2) | 23,846 | 0.028 | 0.016 (0.061) | 0.050 (0.029) | 0.005 (0.018) | 0.032 (0.017) | 0.13 (-0.37, 0.63) |
| Obesity (class 3) | 20,966 | 0.01 | -0.088 (0.105) | 0.046 (0.026) | -0.017 (0.020) | 0.036 (0.020) | N/A |
| Overweight | 57,451 | 0.042 | 0.052 (0.045) | 0.035 (0.010) | 0.009 (0.008) | 0.028 (0.008) | 0.25 (0.04, 0.46) |
| Waist-hip ratio | 57,590 | 0.077 | -0.049 (0.041) | 0.015 (0.010) | -0.010 (0.008) | 0.012 (0.008) | N/A |
| Waist-hip ratio (adjusted for BMI) | 57,388 | 0.065 | -0.115 (0.042) | 0.007 (0.010) | -0.022 (0.008) | 0.005 (0.008) | N/A |
| Waist-hip ratio (men) | 23,387 | 0.004 | -0.081 (0.110) | 0.026 (0.022) | -0.013 (0.018) | 0.022 (0.018) | N/A |
| Waist-hip ratio (women) | 34,203 | -0.011 | -0.023 (0.055) | -0.002 (0.018) | -0.006 (0.013) | -0.001 (0.013) | 0.80 (-1.07, 2.68) |
| Waist circumference | 57,641 | 0.103 | 0.084 (0.041) | 0.091 (0.010) | 0.017 (0.008) | 0.072 (0.008) | 0.19 (0.10, 0.28) |
| Waist circumference (adjusted for BMI) | 57,435 | 0.129 | 0.147 (0.041) | 0.008 (0.011) | 0.029 (0.008) | 0.006 (0.008) | 0.83 (0.60, 1.05) |
| Waist circumference (men) | 23,318 | 0.072 | 0.090 (0.089) | 0.047 (0.025) | 0.019 (0.018) | 0.035 (0.018) | 0.35 (0.01, 0.68) |
| Waist circumference (women) | 34,252 | 0.014 | 0.083 (0.052) | 0.017 (0.018) | 0.021 (0.013) | 0.012 (0.013) | 0.63 (0.25, 1.01) |
| Hip circumference | 57,559 | 0.066 | 0.173 (0.036) | 0.033 (0.011) | 0.039 (0.008) | 0.025 (0.008) | 0.61 (0.48, 0.73) |
| Hip circumference (adjusted for BMI) | 57,359 | 0.070 | 0.292 (0.037) | 0.015 (0.011) | 0.065 (0.008) | 0.011 (0.008) | 0.85 (0.75, 0.96) |
| Hip circumference (men) | 23,376 | 0.102 | 0.193 (0.082) | 0.058 (0.026) | 0.044 (0.018) | 0.041 (0.018) | 0.51 (0.30, 0.73) |
| Hip circumference (women) | 34,183 | 0.046 | 0.146 (0.053) | 0.026 (0.018) | 0.037 (0.013) | 0.019 (0.013) | 0.66 (0.43, 0.89) |
| Age at menarche | 33,518 | 0.022 | 0.021 (0.054) | 0.008 (0.019) | 0.005 (0.013) | 0.006 (0.013) | 0.49 (-0.72, 1.71) |
| Asthma | 57,715 | -0.019 | -0.014 (0.054) | -0.007 (0.010) | -0.002 (0.008) | -0.006 (0.008) | 0.25 (-0.67, 1.16) |

| Adult phenotype | N ^a | Phenotypic correlation with BW | Genetic correlation with BW (SE) | Residual correlation with BW (SE) | Genetic covariance with BW, gcov (SE) | Residual covariance with BW, rcov (SE) | Proportion of covariance explained by genetic variants directly genotyped: gcov/(gcov+rcov) ^b (95% CI) |
|--------------------------|----------------|--------------------------------|----------------------------------|-----------------------------------|---------------------------------------|--|---|
| Ever smoked | 56,960 | 0.042 | -0.011 (0.049) | 0.043 (0.010) | -0.002 (0.008) | 0.035 (0.008) | N/A |
| Former smoker | 50,907 | 0.046 | -0.038 (0.059) | 0.048 (0.011) | -0.006 (0.009) | 0.040 (0.009) | N/A |
| Cigarettes per day | 50,641 | 0.044 | -0.053 (0.052) | 0.049 (0.011) | -0.009 (0.009) | 0.040 (0.009) | N/A |
| Smoking start age | 19,639 | -0.024 | 0.465 (0.239) | -0.054 (0.024) | 0.043 (0.020) | -0.048 (0.020) | N/A |
| NFBC1966 | | | | | | | |
| Triglycerides | 4,954 | -0.065 | -0.448 (0.224) | 0.022 (0.049) | -0.083 (0.038) | 0.018 (0.039) | N/A |
| Total cholesterol | 4,970 | -0.021 | -0.271 (0.137) | 0.080 (0.054) | -0.078 (0.039) | 0.057 (0.038) | N/A |
| High density lipoprotein | 4,972 | 0.020 | 0.101 (0.125) | -0.017 (0.055) | 0.031 (0.039) | -0.011 (0.038) | N/A |
| Low density lipoprotein | 4,970 | -0.013 | -0.223 (0.119) | 0.093 (0.058) | -0.074 (0.039) | 0.062 (0.038) | N/A |
| Fasting glucose | 4,465 | -0.042 | -0.184 (0.173) | 0.006 (0.058) | -0.045 (0.042) | 0.005 (0.042) | N/A |
| Fasting insulin | 4,441 | -0.100 | -0.180 (0.214) | -0.087 (0.056) | -0.036 (0.043) | -0.067 (0.043) | 0.35 (-0.07, 1.28) |

^aThe number of case/control was T2D: 1,488/55,166, CAD: 1,959/55,756, obesity1: 19,820/13,802, obesity2: 4,026/13,802, obesity3: 1,146/13,802, overweight: 37,631/19,820, asthma: 7,131/50,584, ever smoker: 24,774/32,186, and former smoker: 18,721/32,186. The covariance for these traits were on the observed scale.

^bWe have put N/A (not applicable) when the genetic and residual covariances are in the opposite direction and it is not relevant to calculate gcov/(gcov + rcov). Although genetic covariance is subject to high uncertainty, there is a general pattern across both UK Biobank and the NFBC1966 that the phenotypic covariation between low BW and future cardiometabolic risk is at least in part genetically mediated.

Supplementary Table 14. List of 77 reported imprinted regions used in the current analysis.

| Gene | Type | Entrez ID | Chr | Gene Start site (bp, b37) | Gene End site (bp, b37) |
|----------------|-----------------------------|-----------|-----|------------------------------|----------------------------|
| <i>CPA4</i> | GTEX Primary | 51200 | 7 | 129,932,973 | 129,964,020 |
| <i>CST1</i> | GTEX Primary | 1469 | 20 | 23,728,189 | 23,731,574 |
| <i>DIRAS3</i> | GTEX Primary | 9077 | 1 | 68,511,644 | 68,516,460 |
| <i>DLK1</i> | GTEX Primary | 8788 | 14 | 101,193,201 | 101,201,467 |
| <i>FAM50B</i> | GTEX Primary | 26240 | 6 | 3,849,631 | 3,851,551 |
| <i>GRB10</i> | GTEX Primary | 2887 | 7 | 50,657,759 | 50,861,159 |
| <i>H19</i> | GTEX Primary | 283120 | 11 | 2,016,405 | 2,019,065 |
| <i>IGF2</i> | GTEX Primary | 3481 | 11 | 2,150,346 | 2,170,833 |
| <i>IGF2AS</i> | GTEX Primary | 51214 | 11 | 2,161,757 | 2,169,896 |
| <i>INPP5F</i> | GTEX Primary | 22876 | 10 | 121,485,608 | 121,588,659 |
| <i>KCNQ1</i> | GTEX Primary | 3784 | 11 | 2,466,220 | 2,870,340 |
| <i>KIF25</i> | GTEX Primary | 3834 | 6 | 168,418,552 | 168,445,769 |
| <i>L3MBTL1</i> | GTEX Primary | 26013 | 20 | 42,136,319 | 42,170,534 |
| <i>LPAR6</i> | GTEX Primary | 10161 | 13 | 48,985,181 | 49,018,840 |
| <i>MAGEL2</i> | GTEX Primary | 54551 | 15 | 23,888,695 | 23,892,993 |
| <i>MAGI2</i> | GTEX Primary | 9863 | 7 | 77,646,373 | 79,082,890 |
| <i>MEG3</i> | GTEX Primary | 55384 | 14 | 101,292,444 | 101,327,360 |
| <i>MEG8</i> | GTEX Primary | 79104 | 14 | 101,361,106 | 101,373,305 |
| <i>MEST</i> | GTEX Primary | 4232 | 7 | 130,126,045 | 130,146,131 |
| <i>NAP1L5</i> | GTEX Primary | 266812 | 4 | 89,617,065 | 89,619,023 |
| <i>NDN</i> | GTEX Primary | 4692 | 15 | 23,930,553 | 23,932,450 |
| <i>NTM</i> | GTEX Primary | 50863 | 11 | 131,240,370 | 132,206,716 |
| <i>PEG10</i> | GTEX Primary | 23089 | 7 | 94,285,636 | 94,299,006 |
| <i>PEG3</i> | GTEX Primary | 5178 | 19 | 57,321,444 | 57,352,094 |
| <i>PLAGL1</i> | GTEX Primary | 5325 | 6 | 144,261,436 | 144,385,735 |
| <i>PPIEL</i> | GTEX Primary | 728448 | 1 | 39,987,951 | 40,025,370 |
| <i>PPP2R5D</i> | GTEX Primary | 5528 | 6 | 42,952,329 | 42,980,080 |
| <i>PWRN1</i> | GTEX Primary | 791114 | 15 | 24,803,303 | 24,832,926 |
| <i>SGK2</i> | GTEX Primary | 10110 | 20 | 42,187,634 | 42,214,273 |
| <i>SNRPN</i> | GTEX Primary | 6638 | 15 | 25,068,793 | 25,223,729 |
| <i>SNURF</i> | GTEX Primary | 8926 | 15 | 25,200,069 | 25,223,729 |
| <i>SYCE1</i> | GTEX Primary | 93426 | 10 | 135,367,403 | 135,382,876 |
| <i>UBE3A</i> | GTEX Primary | 7337 | 15 | 25,582,395 | 25,684,128 |
| <i>UGT2B4</i> | GTEX Primary | 7363 | 4 | 70,345,882 | 70,361,626 |
| <i>UTS2</i> | GTEX Primary | 10911 | 1 | 7,907,671 | 7,913,551 |
| <i>ZDBF2</i> | GTEX Primary | 57683 | 2 | 207,139,522 | 207,179,148 |
| <i>ZNF331</i> | GTEX Primary | 55422 | 19 | 54,024,176 | 54,083,523 |
| <i>ZNF597</i> | GTEX Primary | 146434 | 16 | 3,486,109 | 3,493,490 |
| <i>ABCA6</i> | GTEX suggestive (S4 unique) | 23460 | 17 | 67,074,846 | 67,138,015 |
| <i>BMP8A</i> | GTEX suggestive (S4 unique) | 353500 | 1 | 39,957,317 | 39,995,541 |
| <i>CHIT1</i> | GTEX suggestive (S4 unique) | 1118 | 1 | 203,185,206 | 203,198,860 |
| <i>DLGAP2</i> | GTEX suggestive (S4 unique) | 9228 | 8 | 1,449,568 | 1,656,642 |
| <i>DZIP1</i> | GTEX suggestive (S4 unique) | 22873 | 13 | 96,230,455 | 96,296,957 |
| <i>EMR1</i> | GTEX suggestive (S4 unique) | 2015 | 19 | 6,887,581 | 6,940,464 |
| <i>FKBP10</i> | GTEX suggestive (S4 unique) | 60681 | 17 | 39,968,961 | 39,979,469 |
| <i>GPR1</i> | GTEX suggestive (S4 unique) | 2825 | 2 | 207,040,041 | 207,082,771 |
| <i>GTSF1</i> | GTEX suggestive (S4 unique) | 121355 | 12 | 54,849,735 | 54,867,386 |
| <i>GUCY1B2</i> | GTEX suggestive (S4 unique) | 2974 | 13 | 51,568,646 | 51,640,293 |
| <i>KIR3DX1</i> | GTEX suggestive (S4 unique) | 90011 | 19 | 55,043,908 | 55,055,195 |
| <i>LRRTM1</i> | GTEX suggestive (S4 unique) | 347730 | 2 | 80,529,002 | 80,531,487 |

| Gene | Type | Entrez ID | Chr | Gene Start site (bp, b37) | Gene End site (bp, b37) |
|-----------------|-----------------------------|-----------|-----|------------------------------|----------------------------|
| <i>MYOM2</i> | GTEX suggestive (S4 unique) | 9172 | 8 | 1,993,157 | 2,093,380 |
| <i>RTL1</i> | GTEX suggestive (S4 unique) | 388015 | 14 | 101,346,991 | 101,351,184 |
| <i>SERPINA6</i> | GTEX suggestive (S4 unique) | 866 | 14 | 94,770,584 | 94,789,688 |
| <i>SLC10A2</i> | GTEX suggestive (S4 unique) | 6555 | 13 | 103,696,347 | 103,719,196 |
| <i>THNSL2</i> | GTEX suggestive (S4 unique) | 55258 | 2 | 88,469,834 | 88,486,146 |
| <i>ANO1</i> | GTEX consistent (S6) | 55107 | 11 | 69,924,407 | 70,035,652 |
| <i>CALCR</i> | GTEX consistent (S6) | 799 | 7 | 93,053,798 | 93,204,042 |
| <i>CDKN1C</i> | GTEX consistent (S6) | 1028 | 11 | 2,904,447 | 2,906,995 |
| <i>COPG2</i> | GTEX consistent (S6) | 26958 | 7 | 130,146,079 | 130,353,598 |
| <i>DCN</i> | GTEX consistent (S6) | 1634 | 12 | 91,539,034 | 91,576,806 |
| <i>DIO3</i> | GTEX consistent (S6) | 1735 | 14 | 102,027,687 | 102,029,789 |
| <i>DLX5</i> | GTEX consistent (S6) | 1749 | 7 | 96,649,701 | 96,654,143 |
| <i>GABRA5</i> | GTEX consistent (S6) | 2558 | 15 | 27,111,865 | 27,194,357 |
| <i>GABRG3</i> | GTEX consistent (S6) | 2567 | 15 | 27,216,428 | 27,778,373 |
| <i>GLIS3</i> | GTEX consistent (S6) | 169792 | 9 | 3,824,127 | 4,300,035 |
| <i>GNAS-AS1</i> | GTEX consistent (S6) | 149775 | 20 | 57,393,972 | 57,425,958 |
| <i>KLF14</i> | GTEX consistent (S6) | 136259 | 7 | 130,417,395 | 130,418,888 |
| <i>MKRN3</i> | GTEX consistent (S6) | 7681 | 15 | 23,810,453 | 23,813,166 |
| <i>NNAT</i> | GTEX consistent (S6) | 4826 | 20 | 36,149,606 | 36,152,090 |
| <i>C15orf2</i> | GTEX consistent (S6) | 23742 | 15 | 24,920,540 | 24,928,593 |
| <i>PON3</i> | GTEX consistent (S6) | 5446 | 7 | 94,989,183 | 95,025,687 |
| <i>RB1</i> | GTEX consistent (S6) | 5925 | 13 | 48,877,882 | 49,056,026 |
| <i>RBP5</i> | GTEX consistent (S6) | 83758 | 12 | 7,276,279 | 7,281,466 |
| <i>SGCE</i> | GTEX consistent (S6) | 8910 | 7 | 94,214,535 | 94,285,521 |
| <i>TH</i> | GTEX consistent (S6) | 7054 | 11 | 2,185,158 | 2,193,035 |
| <i>TSPAN32</i> | GTEX consistent (S6) | 10077 | 11 | 2,323,242 | 2,339,430 |
| <i>TSSC4</i> | GTEX consistent (S6) | 10078 | 11 | 2,423,522 | 2,425,106 |

These 77 regions were highlighted in reference 10: Baran Y *et al. Genome Res* **25**, 927-936 (2015).

Supplementary Table 15. Comparison of variance in BW between individuals heterozygous and homozygous for each of the 59 autosomal BW index SNPs (plus *DLK1*) in 57,715 UK Biobank samples.

| Locus | Index variant | Alleles A/B | Variance | | | Effect | | |
|--------------------|---------------|----------------|----------|-------|-------|--------------------|-------|----------------------|
| | | | AA | AB | BB | β | SE | P-value |
| <i>WNT4-ZBTB40</i> | rs2473248 | T/C | 1.008 | 0.971 | 1.007 | -0.015 | 0.006 | 0.99 |
| <i>ZBTB7B</i> | rs3753639 | T/C | 0.994 | 1.004 | 1.026 | 0.002 | 0.005 | 0.34 |
| <i>FCGR2B</i> | rs72480273 | A/C | 0.995 | 1.008 | 1.012 | 0.008 | 0.005 | 6.4×10^{-2} |
| <i>DTL</i> | rs61830764 | G/A | 0.997 | 0.993 | 1.031 | -0.004 | 0.005 | 0.81 |
| <i>ATAD2B</i> | rs7575873 | A/G | 1.007 | 0.978 | 0.964 | -0.014 | 0.006 | 0.99 |
| <i>EPAS1</i> | rs1374204 | C/T | 0.968 | 0.999 | 1.005 | -0.002 | 0.005 | 0.68 |
| <i>PTH1R</i> | rs2242116 | A/G | 1.003 | 0.987 | 1.013 | -0.010 | 0.005 | 0.98 |
| <i>ADCY5</i> | rs11719201 | C/T | 0.984 | 1.019 | 1.023 | 0.016 | 0.005 | 1.1×10^{-3} |
| <i>CPA3</i> | rs10935733 | T/C | 0.999 | 1.007 | 0.990 | 0.009 | 0.005 | 3.7×10^{-2} |
| <i>CCNL1-LEKR1</i> | rs13322435 | A/G | 1.016 | 0.993 | 0.981 | -0.007 | 0.005 | 0.91 |
| <i>LCORL</i> | rs925098 | G/A | 1.011 | 0.996 | 1.001 | -0.002 | 0.005 | 0.64 |
| <i>HHIP</i> | rs6537307 | A/G | 0.972 | 1.004 | 1.019 | 0.005 | 0.005 | 0.18 |
| <i>5q11.2</i> | rs854037 | A/G | 1.004 | 0.990 | 1.008 | -0.009 | 0.005 | 0.95 |
| <i>EBF1</i> | rs7729301 | G/A | 1.028 | 0.990 | 1.003 | -0.007 | 0.005 | 0.93 |
| <i>CDKAL1</i> | rs35261542 | C/A | 1.004 | 0.998 | 0.969 | 2×10^{-4} | 0.005 | 0.48 |
| <i>HIST1H2BE</i> | rs9379832 | A/G | 1.004 | 0.995 | 0.995 | -0.004 | 0.005 | 0.80 |
| <i>HMGA1</i> | rs7742369 | A/G | 0.995 | 1.005 | 1.044 | 0.002 | 0.005 | 0.36 |
| <i>L3MBTL3</i> | rs1415701 | G/A | 1.006 | 0.990 | 1.006 | -0.009 | 0.005 | 0.96 |
| <i>ESR1</i> | rs1101081 | C/T | 1.003 | 0.999 | 0.979 | 0.001 | 0.005 | 0.39 |
| <i>GNA12</i> | rs798489 | C/T | 1.005 | 0.993 | 0.997 | -0.005 | 0.005 | 0.81 |
| <i>IGF2BP3</i> | rs11765649 | T/C | 1.004 | 0.991 | 1.007 | -0.005 | 0.005 | 0.83 |
| <i>TBX20</i> | rs6959887 | A/G | 0.998 | 1.001 | 0.999 | 0.004 | 0.005 | 0.21 |
| <i>YKT6-GCK</i> | rs138715366 | C/T | 1.001 | 0.888 | 0.657 | -0.070 | 0.019 | 1.00 |
| <i>MLXIPL</i> | rs62466330 | T/C | 0.995 | 1.028 | 1.036 | 0.016 | 0.007 | 1.3×10^{-2} |
| <i>ANK1-NKX6-3</i> | rs13266210 | A/G | 1.007 | 0.984 | 1.009 | -0.013 | 0.005 | 0.99 |
| <i>TRIB1</i> | rs6989280 | G/A | 0.997 | 1.005 | 0.992 | 0.005 | 0.005 | 0.18 |
| <i>SLC45A4</i> | rs12543725 | G/A | 1.009 | 0.994 | 0.999 | -0.009 | 0.005 | 0.97 |
| <i>PTCH1</i> | rs28510415 | A/G | 0.999 | 0.999 | 1.017 | 1×10^{-4} | 0.007 | 0.49 |
| <i>LPAR1</i> | rs2150052 | A/T | 0.980 | 1.002 | 1.015 | 0.002 | 0.005 | 0.37 |
| <i>PHF19</i> | rs7847628 | A/G | 0.984 | 1.002 | 1.001 | 0.003 | 0.005 | 0.30 |
| <i>STRBP</i> | rs700059 | G/A | 0.989 | 1.018 | 0.994 | 0.015 | 0.006 | 4.4×10^{-3} |
| <i>HHEX-IDE</i> | rs61862780 | T/C | 1.005 | 0.996 | 1.003 | -0.005 | 0.005 | 0.82 |
| <i>NT5C2</i> | rs74233809 | T/C | 0.999 | 0.998 | 1.136 | 3×10^{-4} | 0.007 | 0.48 |
| <i>ADRB1</i> | rs7076938 | C/T | 0.992 | 1.003 | 0.998 | 6×10^{-4} | 0.005 | 0.45 |
| <i>PLEKHA1</i> | rs2421016 | C/T | 0.988 | 1.007 | 0.998 | 0.008 | 0.005 | 6.1×10^{-2} |
| <i>INS-IGF2</i> | rs72851023 | C/T | 0.996 | 1.022 | 1.013 | 0.013 | 0.007 | 3.4×10^{-2} |
| <i>MTNR1B</i> | rs10830963 | C/G | 0.994 | 1.008 | 0.997 | 0.005 | 0.005 | 0.18 |
| <i>APOLD1</i> | rs11055034 | C/A | 1.002 | 1.000 | 0.985 | 2×10^{-4} | 0.005 | 0.49 |
| <i>ABCC9</i> | rs139975827 | G/A | 1.006 | 1.000 | 0.982 | 2×10^{-4} | 0.005 | 0.48 |
| <i>ITPR2</i> | rs12823128 | T/C | 1.014 | 0.993 | 0.998 | -0.006 | 0.005 | 0.87 |
| <i>HMGA2</i> | rs1351394 | T/C | 1.003 | 1.001 | 0.993 | 9×10^{-4} | 0.005 | 0.43 |
| <i>IGF1</i> | rs7964361 | G/A | 0.999 | 1.006 | 0.991 | 0.003 | 0.007 | 0.35 |
| <i>LINC00332</i> | rs2324499 | G/C | 1.000 | 0.998 | 1.004 | -0.003 | 0.005 | 0.70 |
| <i>RB1</i> | rs2854355 | A/G | 0.991 | 1.008 | 1.024 | 0.008 | 0.005 | 5.4×10^{-2} |
| <i>RNF219-AS1</i> | rs1819436 | T/C | 1.034 | 1.003 | 0.998 | 5×10^{-4} | 0.006 | 0.47 |
| <i>DLK1</i> | rs6575803 | C/T | 1.002 | 0.991 | 0.987 | -0.007 | 0.006 | 0.87 |
| <i>FES</i> | rs12906125 | G/A | 1.004 | 0.993 | 1.010 | -0.005 | 0.005 | 0.86 |
| <i>IGF1R</i> | rs7402982 | A/G | 0.994 | 0.998 | 1.006 | -0.003 | 0.005 | 0.73 |

| Locus | Index variant | Alleles A/B | Variance | | | Effect | | |
|----------------------|---------------|----------------|----------|-------|-------|--------------------|-------|---------|
| | | | AA | AB | BB | β | SE | P-value |
| <i>GPR139</i> | rs1011939 | G/A | 1.022 | 1.006 | 0.991 | 0.005 | 0.005 | 0.14 |
| <i>CLDN7</i> | rs113086489 | C/T | 0.991 | 1.002 | 1.002 | 0.001 | 0.005 | 0.41 |
| <i>SUZ12P1-CRLF3</i> | rs144843919 | G/A | 1.000 | 0.990 | 0.955 | -0.006 | 0.010 | 0.72 |
| <i>SP6-SP2</i> | rs12942207 | C/T | 1.018 | 1.005 | 0.992 | 0.006 | 0.005 | 0.12 |
| <i>ACTL9</i> | rs61154119 | T/G | 0.998 | 1.007 | 0.974 | 0.003 | 0.006 | 0.30 |
| <i>PEPD</i> | rs10402712 | G/A | 1.005 | 0.993 | 0.998 | -0.004 | 0.005 | 0.78 |
| <i>JAG1</i> | rs6040076 | G/C | 0.989 | 1.003 | 1.003 | 0.004 | 0.005 | 0.22 |
| <i>C20orf203</i> | rs28530618 | A/G | 1.002 | 1.000 | 0.997 | 4×10^{-4} | 0.005 | 0.47 |
| <i>MAFB</i> | rs6016377 | C/T | 0.988 | 0.999 | 1.021 | 2×10^{-4} | 0.005 | 0.48 |
| <i>NRIP1</i> | rs2229742 | G/C | 1.002 | 0.991 | 0.939 | -0.006 | 0.006 | 0.83 |
| <i>KREMEN1</i> | rs134594 | C/T | 1.020 | 1.009 | 0.983 | 0.005 | 0.005 | 0.14 |
| <i>SREBF2</i> | rs62240962 | C/T | 1.003 | 0.985 | 0.983 | -0.008 | 0.007 | 0.89 |

The beta value is a parameter of the Brown-Forsythe test that measures how different the mean absolute deviations (MADs) are from the respective group medians in the heterozygous vs the homozygous group. SE, standard error.

Supplementary Table 16. Parent-of-origin specific analysis at 59 autosomal BW loci (plus *DLK1*) in 4,908 ALSPAC mother-child pairs.

| Locus | SNP | EA/ NEA | Paternal transmission | | | Maternal transmission | | | <i>P</i> _het |
|--------------------|-------------|------------|-----------------------|-------|----------------------|-----------------------|-------|----------------------|----------------------|
| | | | β | SE | <i>P</i> -value | β | SE | <i>P</i> -value | |
| <i>WNT4-ZBTB40</i> | rs2473248 | C/T | 0.019 | 0.042 | 0.65 | 0.189 | 0.076 | 1.3×10^{-2} | 6.1×10^{-2} |
| <i>ZBTB7B</i> | rs3753639 | C/T | 0.012 | 0.034 | 0.72 | 0.078 | 0.045 | 8.5×10^{-2} | 0.34 |
| <i>FCGR2B</i> | rs72480273 | C/A | 0.097 | 0.037 | 8.0×10^{-3} | 0.060 | 0.047 | 0.20 | 0.17 |
| <i>DTL</i> | rs61830764 | A/G | 0.067 | 0.031 | 2.8×10^{-2} | 0.096 | 0.030 | 1.7×10^{-3} | 0.44 |
| <i>ATAD2B</i> | rs7575873 | A/G | 0.080 | 0.041 | 5.3×10^{-2} | 0.086 | 0.072 | 0.23 | 0.36 |
| <i>EPAS1</i> | rs1374204 | T/C | 0.094 | 0.032 | 3.7×10^{-3} | 0.121 | 0.037 | 1.1×10^{-3} | 0.98 |
| <i>PTH1R</i> | rs2242116 | A/G | 0.007 | 0.031 | 0.83 | 0.064 | 0.033 | 5.2×10^{-2} | 9.8×10^{-2} |
| <i>ADCY5</i> | rs11719201 | T/C | 0.137 | 0.034 | 4.5×10^{-3} | 0.062 | 0.035 | 7.5×10^{-2} | 0.11 |
| <i>CPA3</i> | rs10935733 | T/C | 0.052 | 0.029 | 7.7×10^{-2} | 0.013 | 0.031 | 0.67 | 0.19 |
| <i>CCNL1-LEKR1</i> | rs13322435 | A/G | 0.117 | 0.031 | 1.4×10^{-4} | 0.088 | 0.032 | 5.9×10^{-3} | 0.48 |
| <i>LCORL</i> | rs925098 | G/A | 0.128 | 0.033 | 1.1×10^{-4} | 0.063 | 0.042 | 0.13 | 0.18 |
| <i>HHIP</i> | rs6537307 | G/A | 0.080 | 0.030 | 8.1×10^{-3} | 0.071 | 0.030 | 1.9×10^{-2} | 0.79 |
| <i>5q11.2</i> | rs854037 | A/G | 0.078 | 0.037 | 3.5×10^{-2} | 0.081 | 0.043 | 5.8×10^{-2} | 0.90 |
| <i>EBF1</i> | rs7729301 | A/G | -0.006 | 0.034 | 0.86 | -0.028 | 0.041 | 0.50 | 0.67 |
| <i>CDKAL1</i> | rs35261542 | C/A | 0.101 | 0.033 | 2.1×10^{-3} | 0.040 | 0.041 | 0.32 | 8.3×10^{-2} |
| <i>HIST1H2BE</i> | rs9379832 | A/G | 0.034 | 0.033 | 0.30 | 0.037 | 0.038 | 0.33 | 0.83 |
| <i>HMGA1</i> | rs7742369 | G/A | 0.046 | 0.038 | 0.23 | 0.161 | 0.051 | 1.6×10^{-3} | 0.10 |
| <i>L3MBTL3</i> | rs1415701 | G/A | 0.029 | 0.033 | 0.38 | 0.092 | 0.042 | 2.8×10^{-2} | 9.2×10^{-2} |
| <i>ESR1</i> | rs1101081 | C/T | 0.074 | 0.033 | 2.5×10^{-2} | 0.025 | 0.039 | 0.53 | 0.16 |
| <i>GNA12</i> | rs798489 | C/T | 0.011 | 0.033 | 0.74 | 0.043 | 0.040 | 0.29 | 0.35 |
| <i>IGF2BP3</i> | rs11765649 | T/C | -0.013 | 0.033 | 0.69 | 0.054 | 0.036 | 0.14 | 0.14 |
| <i>TBX20</i> | rs6959887 | A/G | 0.035 | 0.032 | 0.27 | 0.035 | 0.033 | 0.29 | 0.94 |
| <i>YKT6-GCK</i> | rs138715366 | C/T | 0.235 | 0.166 | 0.16 | 0.032 | 0.468 | 0.95 | 0.66 |
| <i>MLXIPL</i> | rs62466330 | C/T | 0.107 | 0.053 | 4.5×10^{-2} | 0.166 | 0.094 | 7.8×10^{-2} | 0.87 |
| <i>ANK1-NKX6-3</i> | rs13266210 | A/G | 0.039 | 0.036 | 0.28 | -0.031 | 0.041 | 0.45 | 0.21 |
| <i>TRIB1</i> | rs6989280 | G/A | -0.012 | 0.034 | 0.73 | 0.018 | 0.040 | 0.66 | 0.60 |
| <i>SLC45A4</i> | rs12543725 | G/A | 0.052 | 0.030 | 8.6×10^{-2} | 0.047 | 0.030 | 0.12 | 0.90 |
| <i>PTCH1</i> | rs28510415 | G/A | 0.091 | 0.047 | 5.4×10^{-2} | 0.212 | 0.092 | 2.1×10^{-2} | 0.39 |
| <i>LPAR1</i> | rs2150052 | T/A | -0.001 | 0.030 | 0.97 | 0.092 | 0.030 | 2.0×10^{-3} | 2.4×10^{-3} |
| <i>PHF19</i> | rs7847628 | G/A | 0.038 | 0.032 | 0.23 | 0.004 | 0.037 | 0.92 | 0.28 |
| <i>STRBP</i> | rs700059 | G/A | 0.043 | 0.041 | 0.29 | -0.029 | 0.070 | 0.68 | 0.22 |
| <i>HHEX-IDE</i> | rs61862780 | T/C | 0.085 | 0.030 | 4.7×10^{-3} | 0.050 | 0.030 | 9.6×10^{-2} | 0.32 |
| <i>NT5C2</i> | rs74233809 | C/T | -0.018 | 0.051 | 0.73 | 0.040 | 0.068 | 0.56 | 0.51 |
| <i>ADRB1</i> | rs7076938 | T/C | 0.023 | 0.033 | 0.49 | 0.078 | 0.041 | 5.5×10^{-2} | 0.27 |
| <i>PLEKHA1</i> | rs2421016 | T/C | 0.024 | 0.031 | 0.43 | 0.036 | 0.031 | 0.24 | 0.77 |
| <i>INS-IGF2</i> | rs72851023 | T/C | 0.116 | 0.051 | 2.2×10^{-2} | 0.108 | 0.107 | 0.31 | 0.56 |
| <i>MTNR1B</i> | rs10830963 | G/C | 0.024 | 0.033 | 0.46 | 0.058 | 0.040 | 0.15 | 0.53 |
| <i>APOLD1</i> | rs11055034 | C/A | 0.122 | 0.032 | 1.7×10^{-4} | 0.053 | 0.039 | 0.17 | 3.2×10^{-2} |
| <i>ABCC9</i> | rs139975827 | G/A | 0.045 | 0.031 | 0.15 | 0.017 | 0.032 | 0.59 | 0.46 |
| <i>ITPR2</i> | rs12823128 | T/C | 0.064 | 0.030 | 3.0×10^{-2} | -0.003 | 0.029 | 0.91 | 7.7×10^{-2} |
| <i>HMGA2</i> | rs1351394 | T/C | 0.015 | 0.031 | 0.63 | 0.082 | 0.031 | 8.3×10^{-3} | 8.0×10^{-2} |
| <i>IGF1</i> | rs7964361 | A/G | -0.045 | 0.049 | 0.36 | 0.091 | 0.099 | 0.36 | 0.33 |
| <i>LINC00332</i> | rs2324499 | G/C | 0.012 | 0.032 | 0.70 | 0.041 | 0.034 | 0.22 | 0.70 |
| <i>RB1</i> | rs2854355 | G/A | 0.031 | 0.033 | 0.35 | -0.012 | 0.042 | 0.77 | 0.25 |
| <i>RNF219-AS1</i> | rs1819436 | C/T | 0.038 | 0.042 | 0.36 | 0.136 | 0.079 | 8.4×10^{-2} | 0.39 |
| <i>DLK1</i> | rs6575803 | C/T | 0.104 | 0.045 | 2.2×10^{-2} | 0.205 | 0.091 | 2.5×10^{-2} | 0.40 |
| <i>FES</i> | rs12906125 | G/A | 0.019 | 0.032 | 0.55 | 0.095 | 0.035 | 6.7×10^{-3} | 9.3×10^{-2} |
| <i>IGF1R</i> | rs7402982 | A/G | 0.047 | 0.031 | 0.13 | 0.036 | 0.032 | 0.25 | 0.78 |

| Locus | SNP | EA/ NEA | Paternal transmission | | | Maternal transmission | | | <i>P</i> _het |
|----------------------|-------------|------------|-----------------------|-------|----------------------|-----------------------|-------|----------------------|----------------------|
| | | | β | SE | <i>P</i> -value | β | SE | <i>P</i> -value | |
| <i>GPR139</i> | rs1011939 | G/A | -0.028 | 0.032 | 0.38 | 0.021 | 0.038 | 0.59 | 0.29 |
| <i>CLDN7</i> | rs113086489 | T/C | 0.061 | 0.031 | 4.9×10^{-2} | 0.050 | 0.031 | 0.11 | 0.83 |
| <i>SUZ12P1-CRLF3</i> | rs144843919 | G/A | 0.003 | 0.071 | 0.96 | 0.182 | 0.113 | 0.11 | 0.20 |
| <i>SP6-SP2</i> | rs12942207 | C/T | 0.050 | 0.032 | 0.12 | 0.027 | 0.038 | 0.48 | 0.32 |
| <i>ACTL9</i> | rs61154119 | T/G | 0.056 | 0.038 | 0.14 | 0.164 | 0.062 | 8.7×10^{-3} | 5.6×10^{-2} |
| <i>PEPD</i> | rs10402712 | A/G | 0.077 | 0.033 | 2.1×10^{-2} | 0.001 | 0.034 | 0.98 | 0.10 |
| <i>JAG1</i> | rs6040076 | C/G | 0.028 | 0.031 | 0.36 | 0.001 | 0.031 | 0.97 | 0.53 |
| <i>C20orf203</i> | rs28530618 | A/G | 0.053 | 0.031 | 8.3×10^{-2} | 0.013 | 0.031 | 0.68 | 0.26 |
| <i>MAFB</i> | rs6016377 | T/C | 0.054 | 0.031 | 8.0×10^{-2} | 0.008 | 0.031 | 0.80 | 0.20 |
| <i>NRIP1</i> | rs2229742 | G/C | 0.020 | 0.046 | 0.66 | 0.030 | 0.095 | 0.75 | 0.89 |
| <i>KREMEN1</i> | rs134594 | C/T | 0.024 | 0.031 | 0.43 | 0.052 | 0.035 | 0.13 | 0.75 |
| <i>SREBF2</i> | rs62240962 | C/T | -0.004 | 0.052 | 0.94 | 0.088 | 0.065 | 0.18 | 0.40 |

EA, effect allele; NEA, non effect allele; SE, standard error; *P*_het, *P*-value for difference in BW between AB heterozygote offspring inheriting allele A from mother and B from father, and those inheriting A from father and B from mother.

Supplementary Table 17. Association of BW signals with various adult metabolic and anthropometric traits. (GWAS look-ups)

Provided in a separate Excel file.

Supplementary Table 18. Effect of transmitted and untransmitted maternal haplotype scores on BW in 5,201 ALSPAC mother-child pairs.

| Haplotype score | Height | | | Systolic blood pressure | | | Type 2 diabetes | | |
|-----------------|---------|-------|-----------------------|-------------------------|------|-----------------|-----------------|------|-----------------|
| | β | SE | <i>P</i> -value | β | SE | <i>P</i> -value | β | SE | <i>P</i> -value |
| M1 (C1) | 0.128 | 0.013 | 1.6×10^{-21} | -6.82 | 3.36 | 0.04 | -0.44 | 1.73 | 0.80 |
| M2 | 0.010 | 0.012 | 0.39 | -6.40 | 2.63 | 0.02 | 1.66 | 1.42 | 0.24 |
| C2 | 0.066 | 0.012 | 1.2×10^{-7} | 0.14 | 2.62 | 0.96 | -2.94 | 1.47 | 0.045 |

Beta values are in grams per weighted trait-raising allele. SE, standard error.

M1(C1), maternal transmitted haplotype score; M2, Maternal untransmitted haplotype score; C2, paternal transmitted haplotype score.

Supplementary Table 19. Reciprocal approximate conditional analyses at *YKT6-GCK* variants associated with birth weight (BW) or fasting glucose (FG), in European ancestry meta-analysis of up to 143,677 individuals.

| Variant (chr:position) | Type | EA/ NEA | EAF (Eur) | BW SNP conditioned on FG SNP | | | | FG SNP conditioned on BW SNP | | | |
|-----------------------------|---|------------|--------------|------------------------------|-------------------|---|--------------------------|------------------------------|-------------------|----------------------------|--------------------------|
| | | | | Uncondit ioned <i>P</i> | Conditioned on | Variant type | Conditio ned <i>P</i> | Variant | Conditioned on | Uncondit ioned <i>P</i> | Conditio ned <i>P</i> |
| rs138715366 (7:44246271) | BW lead SNP (at <i>YKT6</i> intron 1) | T/C | 0.0089 | 7.2x10 ⁻²⁶ | unconditioned | - | - | - | - | - | - |
| | | | | | rs878521 | <i>GCK</i> FG primary (ENGAGE ¹) | 1.3x10 ⁻²⁵ | rs878521 | rs138715366 | 0.12 | 0.28 |
| | | | | | rs10259649 | <i>GCK</i> FG secondary (ENGAGE ¹) | 7.7x10 ⁻²⁶ | rs10259649 | rs138715366 | 0.52 | 0.63 |
| | | | | | rs4607517 | <i>GCK</i> FG (MAGIC ²) | 2.6x10 ⁻²⁵ | rs4607517 | rs138715366 | 3.2x10 ⁻⁴ | 1.2x10 ⁻³ |
| | | | | | rs10278336 | <i>GCK</i> T2D (DIAGRAM ³) | 1.8x10 ⁻²⁵ | rs10278336 | rs138715366 | 0.057 | 0.19 |
| rs78412508 (7:44223858) | BW second strongest SNP (at <i>GCK</i> intron 3) | A/G | 0.0095 | 8.9x10 ⁻²⁴ | unconditioned | - | - | - | - | - | - |
| | | | | | rs878521 | <i>GCK</i> FG primary (ENGAGE ¹) | 1.6x10 ⁻²³ | rs878521 | rs78412508 | 0.12 | 0.27 |
| | | | | | rs10259649 | <i>GCK</i> FG secondary (ENGAGE ¹) | 9.7x10 ⁻²⁴ | rs10259649 | rs78412508 | 0.52 | 0.66 |
| | | | | | rs4607517 | <i>GCK</i> FG (MAGIC ²) | 3.1x10 ⁻²³ | rs4607517 | rs78412508 | 3.2x10 ⁻⁴ | 1.2x10 ⁻³ |
| | | | | | rs10278336 | <i>GCK</i> T2D (DIAGRAM ³) | 2.2x10 ⁻²³ | rs10278336 | rs78412508 | 0.057 | 0.18 |


EA, effect allele; NEA, non effect allele; EAF, effect allele frequency; Eur, European; BW, birth weight; FG, fasting glucose; T2D, type 2 diabetes.

1. Horikoshi M *et al.* PLoS Genet 11, e1005230 (2015).

2. Dupuis J *et al.* Nat Genet 42, 105-116 (2010).


3. Morris AP *et al.* Nat Genet 44, 981-990 (2012).

Supplementary Figure 1. BW association enrichment for DNaseI hypersensitive (DHS) sites in 128 cell types and four genic annotations. a, Each cell type DHS annotation and four gene-based annotations were individually tested for enrichment using the Bayes Factors for all variants in the 62 credible sets. The effect estimate (filled circle) and 95% CI (horizontal line) are plotted on the x axis. The 128 cell types (listed on the y axis; see Supplementary Table 8 for details) are categorised according to the description fields from ENCODE (stem cell, neonatal, foetal, embryonic and adult groups). We considered an annotation enriched if the 95% CI did not overlap zero (highlighted in red). **b,** Estimated effect of enrichment (filled circle) and 95% CI (horizontal line) for each categorical field, tested in a joint model, are plotted (see Methods).




Supplementary Figure 2. Estimates of phenotypic, genetic and residual correlations (a) and estimates of genetic and residual covariance (b) between birth weight and adult metabolic or anthropometric traits in UK Biobank ‘white British’ samples (up to N=57,715). See details in Supplementary Table 13.

a



b



Full grants and funding supports for contributing studies.

1958 British Birth Cohort acknowledges use of phenotype and genotype data from the British 1958 Birth Cohort DNA collection, funded by the Medical Research Council grant G0000934 and the Wellcome Trust grant 068545/Z/02. Genotyping for the B58C-WTCCC subset was funded by the Wellcome Trust grant 076113/B/04/Z. The B58C-T1DGC genotyping utilised resources provided by the Type 1 Diabetes Genetics Consortium, a collaborative clinical study sponsored by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), National Institute of Allergy and Infectious Diseases (NIAID), National Human Genome Research Institute (NHGRI), National Institute of Child Health and Human Development (NICHD), and Juvenile Diabetes Research Foundation International (JDRF) and supported by U01 DK062418. B58C-T1DGC GWAS data were deposited by the Diabetes and Inflammation Laboratory, Cambridge Institute for Medical Research (CIMR), University of Cambridge, which is funded by Juvenile Diabetes Research Foundation International, the Wellcome Trust and the National Institute for Health Research Cambridge Biomedical Research Centre; the CIMR is in receipt of a Wellcome Trust Strategic Award (079895). The B58C-GABRIEL genotyping was supported by a contract from the European Commission Framework Programme 6 (018996) and grants from the French Ministry of Research.

ALSPAC is supported by the UK Medical Research Council and the Wellcome Trust (Grant ref: 102215/2/13/2), and the University of Bristol provide core support for ALSPAC.

Barcelona Supercomputing Center This work has been sponsored by the grant SEV-2011-00067 of Severo Ochoa Program, awarded by the Spanish Government. This work was supported by an EFSD/Lilly research fellowship.

CHOP was financially supported by an Institute Development Award from the Children's Hospital of Philadelphia, a Research Development Award from the Cotswold Foundation, NIH grant R01 HD056465 and the Daniel B. Burke Endowed Chair for Diabetes Research (SFAG).

CLHNS was supported by National Institutes of Health grants DK078150, TW05596 and HL085144 and pilot funds from RR20649, ES10126, and DK056350.

The **CoLaus** study was and is supported by research grants from GlaxoSmithKline, the Faculty of Biology and Medicine of Lausanne, and the Swiss National Science Foundation (grants 33CSCO-122661, 33CS30-139468 and 33CS30-148401).

COPSACs greatly acknowledge the private and public research funding allocated to COPSAC and listed on www.copsac.com, with special thanks to The Lundbeck Foundation; Ministry of Health; Danish Council for Strategic Research; The Danish Council for Independent Research and The Capital Region Research Foundation as core supporters. The funding agencies did not have any influence on study design, data collection and analysis, decision to publish or preparation of the manuscript. No pharmaceutical company was involved in the study.

DNBC was established with the support of a major grant from the Danish National Research Foundation. Additional support for the DNBC has been obtained from the Danish Pharmacists' Fund, the Egmont Foundation, the March of Dimes Birth Defects Foundation, the Augustinus Foundation and the Health Fund of the Danish Health Insurance Societies. The generation of GWAS genotype data for the DNBC samples was carried out within the GENEVA consortium with funding provided through the NIH Genes, Environment and Health Initiative (GEI) (U01HG004423, U01HG004438, U01HG004446).

EPIC-Norfolk is supported by Cancer Research UK (SP2024-0201 and SP2024-0204) and the Medical Research Council (G9502233).

The **ERF** study as a part of EUROSPAN (European Special Populations Research Network) was supported by European Commission FP6 STRP grant number 018947 (LSHG-CT-2006-01947) and also received funding from the European Community's Seventh Framework Programme (FP7/2007-2013)/grant agreement HEALTH-F4-2007-201413 by the European Commission under the programme "Quality of Life and Management of the Living Resources" of 5th Framework Programme (no. QLG2-CT-2002-01254). The ERF study was further supported by ENGAGE consortium and Center for Medical Systems Biology (CMSB). High-throughput analysis of the ERF data was supported by joint grant from Netherlands Organisation for Scientific Research and the Russian Foundation for Basic Research (NWO-RFBR 047.017.043).

The **Fenland** Study is funded by the Medical Research Council (MC_U106179471) and Wellcome Trust.

The general design of **Generation R** Study is made possible by financial support from the Erasmus Medical Center, Rotterdam, the Erasmus University Rotterdam, the Netherlands Organization for Health Research and Development (ZonMw), the Netherlands Organisation for Scientific Research (NWO), the Ministry of Health, Welfare and Sport and the Ministry of Youth and Families.

GOYA study was conducted as part of the activities of the Gene-diet Interactions in Obesity project (GENDINOBO, www.gendinob.dk) and the MRC Integrative Epidemiology Unit at the University of Bristol (MC_UU_12013/1-9).

HAPO was supported by US National Institutes of Health (NIH) grants (HD34242, HD34243, HG004415, and DK099820) and by the American Diabetes Association.

HBSC received financial support from the Academy of Finland (project grants 120315, 129287, 218029 and 267561).

Hong Kong HAPO follow-up study was supported by Research Grant Council General Research Fund (CU473408, CU471713) and CUHK Faculty of Medicine Outstanding Fellowship.

INMA study was funded by grants from Instituto de Salud Carlos III (Red INMA G03/176 and CB06/02/0041), FIS-FEDER 03/1615, 04/1509, 04/1112, 04/1931, 05/1079, 05/1052, 06/1213, 07/0314, 09/02647, 11/01007, 11/02591, 13/02032, 13/1944, PI041436, PI081151, 11/00178, 97/0588, 00/0021-2, PI061756, PS0901958, 14/00891, and 14/01687, Spanish Ministry of Science and Innovation (SAF2008-00357), European Commission (ENGAGE project and grant agreement HEALTH-F4-2007-201413), Fundació La Marató de TV3, Generalitat de Catalunya-CIRIT 1999SGR 00241 and Conselleria de Sanitat Generalitat Valenciana. Part of the DNA extractions and genotyping was performed at the Spanish National Genotyping Centre (CEGEN-Barcelona).

Inter99 The Novo Nordisk Foundation Center for Basic Metabolic Research is an independent Research Center at the University of Copenhagen partially funded by an unrestricted donation from the Novo Nordisk Foundation (www.metabol.ku.dk). The study was financially supported by research grants from the Danish Research Council, the Danish Centre for Health Technology Assessment, Novo Nordisk Inc., Research Foundation of Copenhagen County, Ministry of Internal Affairs and Health, the Danish Heart Foundation, the Danish Pharmaceutical Association, the Augustinus Foundation, the Ib Henriksen Foundation, the Becket Foundation, and the Danish Diabetes Association.

Leipzig-Childhood-Obesity was supported by grants from Integrated Research and Treatment Center (IFB) Adiposity Diseases 01EO1001, from the German Research Foundation for the Collaborative Research Center CRC1085 "Atherobesity" project C05, and by the European Commission (Beta-JUDO n° 279153).

The **NEO** study is supported by the participating Departments, the Division and the Board of Directors of the Leiden University Medical Center, and by the Leiden University, Research Profile Area Vascular and Regenerative Medicine.

The Northern Finland Birth Cohort (NFBC) Research program, including the study of the NFBC 1966 and 1986 studies, received financial support from Academy of Finland (1114194, 24300796), NHLBI grant 5R01HL087679 through the STAMPEED program (1R1MH083268-01), ENGAGE project and grant agreement HEALTH-F4-2007-201413, the Medical Research Council (grant G0500539, centre grant G0600705, PrevMetSyn), and the Wellcome Trust (project grant GR069224), UK. The program is currently being funded by the H2020-633595 DynaHEALTH action and Academy of Finland EGEA-project.

NTR obtained funding from the Netherlands Organization for Scientific Research (NWO: MagW/ZonMW grants 916-76-125, 904-61-090, 985-10-002, 904-61-193, 480-04-004, 400-05-717, 481-08-011, 451-04-034, 463-06-001, 912-10-020, Addiction- 31160008 Middelgroot-911-09-032, Spinozapremie 56-464-14192), Center for Medical Systems Biology (CMSB, NWO Genomics), NBIC/BioAssist/RK(2008.024), Biobanking and Biomolecular Resources Research Infrastructure (BBMRI-NL), the VU University's Institute for Health and Care Research (EMGO+) and Neuroscience Campus Amsterdam (NCA), the European Science Foundation (ESF, EU/QLRT-2001-01254), the European Community's Seventh Framework Program (FP7/2007- 2013), ENGAGE (HEALTH-F4-2007-201413); the European Research Council (ERC Advanced, 230374), Rutgers University Cell and DNA Repository (National Institute of Mental Health (NIMH) U24 MH068457-06), and the National Institutes of Health (NIH, R01D0042157-01A; 1RC2MH089995-01; R01DK092127-01). Part of the genotyping and analyses were funded by the Genetic Association Information Network (GAIN) of the Foundation for the US National Institutes of Health, and by grants from GAIN and the NIMH (MH081802).

ORCADES was supported by the Chief Scientist Office of the Scottish Government (CZB/4/276 and CZB/4/710), the Royal Society, the MRC Human Genetics Unit, Arthritis Research UK and the European Union framework program 6 EUROSPAN project (contract no. LSHG-CT-2006-018947).

The **PANIC** study has been financially supported by grants from Ministry of Social Affairs and Health of Finland, Ministry of Education and Culture of Finland, Finnish Innovation Fund Sitra, Social Insurance Institution of Finland, Finnish Cultural Foundation, Juho Vainio Foundation, Foundation for Paediatric Research, Paavo Nurmi Foundation, Paulo Foundation, Diabetes Research Foundation, Research Committee of the Kuopio University Hospital Catchment Area (State Research Funding), Kuopio University Hospital (EVO funding number 5031343) and the city of Kuopio.

RAINE study was supported by the National Health and Medical Research Council of Australia (grant numbers 572613, 403981 and 003209) and the Canadian Institutes of Health Research (grant number MOP-82893).

SCORM was funded by National Medical Research Foundation, Singapore (NMRC/0975/2005) and Biomedical Research Council, Singapore (BMRC 06/1/21/19/466).

SORBS was supported by grants from the German Research Council (DFG - SFB 1052 "Obesity mechanisms"; A01, B01, B03, C01 and SPP 1629 TO 718/2-1), from the German Diabetes Association and from the DHFD (Diabetes Hilfs- und Forschungsfonds Deutschland).

STRIP was supported by Academy of Finland (grants 206374, 251360 and 276861); Juho Vainio Foundation; Finnish Cardiac Research Foundation; Finnish Cultural Foundation; Finnish Ministry of Education and Culture; Sigrid Juselius Foundation; Yrjö Jahnsson Foundation; C.G. Sundell Foundation;

Special Governmental Grants for Health Sciences Research, Turku University Hospital; Foundation for Pediatric Research; and Turku University Foundation.

TEENAGE study has been co-financed by the European Union (European Social Fund (ESF)) and Greek national funds through the Operational Program “Education and Lifelong Learning” of the National Strategic Reference Framework (NSRF)—Research Funding Program: Heracleitus II. Investing in knowledge society through the European Social Fund. This work was funded by the Wellcome Trust (098051).

TDCOB received supports from the Danish Innovation Foundation and The Region Zealand Health and Medical Research Foundation. This study was partly supported by grants from the Program Committee for Individuals, Disease and Society of the Danish Innovation Foundation (grant numbers: 0603-00484B and 0603-00457B) and was part of the research activities in The Impact of our Genomes on Individual Treatment Response in Obese Children (TARGET, www.target.ku.dk) and the Indo-Danish bi-lateral project, Genetics and Systems Biology of Childhood Obesity in India and Denmark (BioChild, www.biochild.ku.dk).

The Novo Nordisk Foundation Center for Basic Metabolic Research, an independent Research Center at the University of Copenhagen, partially funded by an unrestricted donation from the Novo Nordisk Foundation.

University of Oxford, UK, was funded by the British Heart Foundation (grant code SP/13/2/30111), the European Commission (ENGAGE: HEALTH-F4-2007-201413), Medical Research Council (G0601261), and the Wellcome Trust (090532, 098381).

The **Young Finns Study** has been financially supported by the Academy of Finland: grants 286284 (T.L.), 134309 (Eye), 126925, 121584, 124282, 129378 (Salve), 117787 (Gendi), and 41071 (Skidi); the Social Insurance Institution of Finland; Kuopio, Tampere and Turku University Hospital Medical Funds (grant X51001 for T.L.); Juho Vainio Foundation; Paavo Nurmi Foundation; Finnish Foundation of Cardiovascular Research (T.L.); Finnish Cultural Foundation; Tampere Tuberculosis Foundation (T.L.); Emil Aaltonen Foundation (T.L.); and Yrjö Jahnsson Foundation (T.L.).

Grants and funding supports for CHARGE Consortium Hematology Working Group

AGES Funding for Age, Gene/Environment Susceptibility Reykjavik Study (AGES) was made possible by NIA/NIH contract AG000932-2 (2009) Characterization of Normal Genomic Variability. AGES is funded by NIH contract N01-AG-12100, the NIA Intramural Research Program, Hjartavernd (the Icelandic Heart Association) and the Althingi (the Icelandic Parliament).

ARIC The Atherosclerosis Risk in the Communities Study is carried out as a collaborative study supported by National Heart, Lung, and Blood Institute contracts (HHSN268201100005C, HHSN268201100006C, HHSN2682 01100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C and HHSN268 201100012C), R01HL087641, R01HL59367 and R01HL086694; National Human Genome Research Institute contract (U01HG 004402) and National Institutes of Health contract (HHSN 268200625226C). Infrastructure was partly supported by grant number UL1RR025005, a component of the National Institutes of Health and NIH Roadmap for Medical Research. ARIC is also supported by University of North Carolina at Chapel Hill (funded by N01-HC-55015), Baylor Medical College (N01- HC-55016), University of Mississippi Medical Center (N01-HC- 55021), University of Minnesota (N01-HC-55019), Johns Hopkins University (N01-HC-55020), University of Texas Houston (N01-HC- 55017) and University of North Carolina (N01-HC-55018).

CHS Cardiovascular Health Study research was supported by NHLBI contracts HHSN268201200036C, HHSN268200800007C, N01HC55222, N01HC85079, N01HC85080, N01HC85081, N01HC85082, N01HC85083, N01HC85086; and NHLBI grants U01HL080295, R01HL087652, R01HL105756, R01HL103612, and R01HL120393 with additional contribution from the National Institute of Neurological Disorders and Stroke (NINDS). Additional support was provided through R01AG023629 from the National Institute on Aging (NIA). A full list of principal CHS investigators and institutions can be found at CHS-NHLBI.org. The provision of genotyping data was supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR000124, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center.

FHS The National Heart, Lung, and Blood Institute's Framingham Heart Study is a joint project of the National Institutes of Health and Boston University School of Medicine and was supported by the National Heart, Lung, and Blood Institute's Framingham Heart Study (contract No. N01-HC-25195) and its contract with Affymetrix for genotyping services (contract No. N02-HL-6-4278). Analyses reflect the efforts and resource development from the Framingham Heart Study investigators participating in the SNP Health Association Resource (SHARe) project. A portion of this research was conducted using the Linux Cluster for Genetic Analysis (LinGA-II) funded by the Robert Dawson Evans Endowment of the Department of Medicine at the Boston University School of Medicine and Boston Medical Center.

Rotterdam Study The generation and management of GWAS genotype data for the Rotterdam Study (RS I, RS II, RS III) was executed by the Human Genotyping Facility of the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, Rotterdam, The Netherlands. The GWAS datasets are supported by the Netherlands Organisation of Scientific Research NWO Investments (nr. 175.010.2005.011, 911-03-012), the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, the Research Institute for Diseases in the Elderly (014-93-015; RIDE2), the Netherlands Genomics Initiative (NGI)/Netherlands Organisation for Scientific Research (NWO) Netherlands Consortium for Healthy Aging (NCHA), project nr. 050-060-810. The Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, Rotterdam, Netherlands Organization for the Health Research and Development (ZonMw), the Research Institute for Diseases in the Elderly (RIDE), the Ministry of Education, Culture and Science, the Ministry for Health, Welfare and Sports, the European Commission (DG XII), and the Municipality of Rotterdam.

The InChianti Study was supported as a 'targeted project' (ICS 110.1RS97.71) by the Italian Ministry of Health, by the US National Institute on Aging (contracts N01-AG-916413, N01- AG-821336, 263 MD 9164 13 and 263 MD 821336) and in part by the Intramural Research Program, National Institute on Aging and National Institutes of Health, USA.

Grants and funding supports for individuals

R.N.B. is funded by the Wellcome Trust and Royal Society (grant: 104150/Z/14/Z). N.M.W. is supported by a National Health and Medical Research Council Early Career Fellowship (APP1104818), a University of Queensland Early Career Researcher Grant (2014002959) and a University of Western Australia - University of Queensland Bilateral Research Collaboration Award (2014001711). B.F. is supported by an Oak Foundation fellowship and by a grant from the Novo Nordisk Foundation (12955). N.R.v.Z. has received salary support from the Innovative Medicines Initiative and the Juvenile Diabetes Research Foundation. T.S.A. was funded by the Gene-Diet Interactions in Obesity (GENDINO) project and acknowledges the same. N.V-T. is funded by a pre-doctoral grant from the Agència de Gestió d'Ajuts

Universitat de Recerca (2015 FI_B 00636), Generalitat de Catalunya. V.L. and I.P. were funded in part through the European Union's Seventh Framework Programme (FP7/2007–2013), ENGAGE project, grant agreement HEALTH-F4-2007-201413. L.A.D received a College of Medical, Veterinary and Life Sciences Medical Research Council Doctoral Training Grant Scholarship (G0900185-1/1). J.M.M. was supported by Sara Borrell Fellowship from the Instituto Carlos III and Beatriu de Pinós fellowship from the Agency for Management of University and Research Grants (AGAUR). N.R. is funded by the Medical Research Council, UK (MR/K011480/1). S.K.G. is supported by HL122684. S.E.J. is funded by the Medical Research Council (grant: MR/M005070/1). M.A.T. is supported by a Wellcome Trust Institutional Strategic Support Award (WT097835MF). J.T. is funded by a Diabetes Research and Wellness Foundation Fellowship. A.R.W. and T.M.F. are supported by the European Research Council grant: 323195 SZ-245 50371-GLUCOSEGENES-FP7-IDEAS-ERC. I.P. is supported by Elsie Widdowson fellowship from the Imperial College London, UK. S.B-G. was FI-DGR Fellowship from FIDGR 2013 from Agència de Gestió d'Ajuts Universitaris i de Recerca (AGAUR, Generalitat de Catalunya). G.H. works within the MRC Integrative Epidemiology Unit at the University of Bristol (MC_UU_12013/1-9). G.M. is supported by core ALSPAC funding Grant ref: 102215/2/13/2). S.M.M. was partly funded by a Research Councils UK Fellowship. E.Z. is supported by the Wellcome Trust (098051). H.N.K. is awarded EU-FP7 Marie Curie Actions - Career Integration Grant (CIG-293511). S.S. and M-R.J. have received funding from the European Union's Horizon 2020 research and innovation programme [under grant agreement No 633595] for the DynaHEALTH action. A.T.H. is a Wellcome Trust Senior Investigator (098395/Z/12/Z) and an NIHR Senior Investigator (NF-SI-0611-10219). P.V. received funding from the Swiss National Science Foundation (grant no: 33CSCO-122661 and FN 33CSCO-139468), GlaxoSmithKline and the Faculty of Biology and Medicine of Lausanne, Switzerland, to conduct the CoLaus study. D.M-K. was supported by Dutch Science Organization (ZonMW-VENI Grant 916.14.023). C.Pisinger is funded by the Health Foundation (Helsefonden). C.Power at UCL Institute of Child Health, with support from the National Institute for Health Research Biomedical Research Centre at Great Ormond Street Hospital for Children NHS Foundation Trust and University College London. B.R.W. is funded by British Heart Foundation and the Wellcome Trust. V.W.V.J. received an additional grant from the Netherlands Organization for Health Research and Development (NWO, ZonMw-VIDI 016.136.361) and a European Research Council Consolidator Grant (ERC-2014-CoG-648916). S.F.A.G. is funded by the Daniel B. Burke Endowed Chair for Diabetes Research. D.A.L. works within the MRC Integrative Epidemiology Unit at the University of Bristol (MC_UU_12013/5). G.D.S. works within the MRC Integrative Epidemiology Unit at the University of Bristol (MC_UU_12013/1). A.P.M. is a Wellcome Trust Senior Research Fellow (grant number WT098017) and acknowledges funding under WT090532 and WT064890. J.F.F. has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 633595 for the DynaHEALTH action. N.J.T. works within the MRC Integrative Epidemiology Unit at the University of Bristol (MC_UU_12013/3). D.M.E. is funded by an Australian Research Council Future Fellowship (FT130101709) and works within the MRC Integrative Epidemiology Unit at the University of Bristol (MC_UU_12013/4). M.I.M. is a Wellcome Trust Senior Investigator (grant number 098381) and a NIHR Senior Investigator. R.M.F. is a Sir Henry Dale Fellow (Wellcome Trust and Royal Society grant: 104150/Z/14/Z).

Study acknowledgements

ALSPAC We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. GWAS data were generated by Sample Logistics and Genotyping Facilities at the Wellcome Trust Sanger Institute and LabCorp (Laboratory Corporation of America) using support from 23andMe.

Barcelona Supercomputing Center acknowledges PRACE for awarding us access to MareNostrum supercomputer, based in Spain at Barcelona.

CHOP authors thank the network of primary care clinicians and the patients and families for their contribution to this project and to clinical research facilitated by the Pediatric Research Consortium (PeRC) at The Children's Hospital of Philadelphia. R. Chiavacci, E. Dabaghyan, A. (Hope) Thomas, K. Harden, A. Hill, C. Johnson-Honesty, C. Drummond, S. Harrison, F. Salley, C. Gibbons, K. Lilliston, C. Kim, E. Frackelton, G. Otieno, K. Thomas, C. Hou, K. Thomas and M.L. Garris provided expert assistance with genotyping and/or data collection and management. The authors would also like to thank S. Kristinsson, L.A. Hermannsson and A. Krisbjörnsson of Raförnninn ehf for extensive software design and contributions.

CLHNS We thank the USC-Office of Population Studies Foundation, Inc. research and data collection teams and the study participants who generously provided their time for this study.

COPSAC2000, COPSAC2010 and COPSAC-REGISTRY gratefully express our gratitude to the participants of the COPSAC2000, COPSAC2010 and COPSAC-REGISTRY cohort study for all their support and commitment. We also acknowledge and appreciate the unique efforts of the COPSAC research team.

DNBC We are very grateful to all the women and children taking part in the DNBC.

EPIC-Norfolk We thank all EPIC-Norfolk study participants and staff for their contribution to the study.

ERF We are grateful to all study participants and their relatives, general practitioners and neurologists for their contributions and to P. Veraart for her help in genealogy, J. Vergeer for the supervision of the laboratory work and P. Snijders for his help in data collection.

Fenland We are grateful to all the volunteers for their time and help, and to the General Practitioners and practice staff for assistance with recruitment. We thank the Fenland Study Investigators, Fenland Study Co-ordination team and the Epidemiology Field, Data and Laboratory teams.

The Generation R Study is conducted by the Erasmus Medical Center in close collaboration with the School of Law and Faculty of Social Sciences of the Erasmus University Rotterdam, the Municipal Health Service Rotterdam area, Rotterdam, the Rotterdam Homecare Foundation, Rotterdam and the Stichting Trombosedienst & Artsenlaboratorium Rijnmond (STAR-MDC), Rotterdam. We gratefully acknowledge the contribution of children and parents, general practitioners, hospitals, midwives and pharmacies in Rotterdam. The study protocol was approved by the Medical Ethical Committee of the Erasmus Medical Center, Rotterdam. Written informed consent was obtained for all participants. The generation and management of GWAS genotype data was executed by the Human Genotyping Facility of the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, Rotterdam, the Netherlands. We thank Pascal Arp, Mila Jhamai, Marijn Verkerk, Lizbeth Herrera and Marjolein Peters for their help in creating, managing and QC of the GWAS database. Also Karol Estrada and Carolina Medina-Gomez for their support in creation and analysis of imputed data. We would like to thank Anis Abuseiris, Karol Estrada, Dr. Tobias A. Knoch, and Rob de Graaf as well as their institutions Biophysical Genomics, Erasmus MC Rotterdam, The Netherlands, and especially the national German MediGRID and Services@MediGRID part of the German D-Grid, both funded by the German Bundesministerium fuer Forschung und Technology under grants #01 AK 803 A-H and # 01 IG 07015 G for access to their grid resources.

GINIplus&LISApplus We thank all families for participation in the studies and the LISApplus and GINIplus study teams for their excellent work. The GINIplus study team wishes to acknowledge the following: Helmholtz Zentrum Muenchen - German Research Center for Environmental Health, Institute of Epidemiology I, Munich (Heinrich J, Brüske I, Schulz H, Standl M, Thiering E, Tiesler CMT, Chen C-M, Schnappinger M); Department of Pediatrics, Marien-Hospital, Wesel (Berdel D, von Berg A); Department

of Pediatrics, Ludwig Maximilians University, Munich (Koletzko S); Department of Pediatrics, Technical University, Munich (Bauer CP, Hoffmann U); IUF – Leibniz Research Institute for Environmental Medicine, Düsseldorf (Krämer U, Link E, Cramer C). The LISApplus study team wishes to acknowledge the following: Helmholtz Zentrum München - German Research Center for Environment and Health, Institute of Epidemiology I, Neuherberg (Heinrich J, Brüske I, Schulz H, Standl M, Thiering E, Tiesler CMT, Chen C-M, Schnappinger M); Department of Pediatrics, Marien-Hospital, Wesel (von Berg A); Bad Honnef (Schaaf B); UFZ-Centre for Environmental Research Leipzig-Halle, Department of Environmental Immunology (Lehmann I); IUF – Leibniz Research Institute for Environmental Medicine, Düsseldorf (Krämer U); Department of Pediatrics, Technical University, Munich (Bauer CP, Hoffman U).

GOYA We thank all the participants of the study.

The Helsinki Birth Cohort Study (HBCS/HBCS 1934-44) thanks Professor David Barker and Tom Forsen. The DNA extraction, sample quality control, biobank up-keep and aliquotting were performed at the National Institute for Health and Welfare, Helsinki, Finland.

INMA authors are grateful to Silvia Fochs, Anna Sánchez, Maribel López, Nuria Pey, Muriel Ferrer, Amparo Quiles, Sandra Pérez, Gemma León, Elena Romero, Maria Andreu, Nati Galiana, Maria Dolores Climent and Amparo Cases for their assistance in contacting the families and administering the questionnaires. The authors would particularly like to thank all the participants for their generous collaboration. A full roster of the INMA Project Investigators can be found at http://www.proyectoinma.org/presentacion-inma/listado-investigadores/en_listado-investigadores.html.

Inter99 study We thank all participants and our committed staff.

Leipzig-Childhood-Obesity are grateful to all the patients and families for contributing to the study. We highly appreciate the support of the Obesity Team and Auxo Team of the Leipzig University Children's Hospital for management of the patients and to the Pediatric Research Center Lab Team for support with DNA banking.

The authors of the **NEO study** thank all individuals who participated in the Netherlands Epidemiology in Obesity study, all participating general practitioners for inviting eligible participants and all research nurses for collection of the data. We thank the NEO study group and Ingeborg de Jonge for the coordination and data management of the NEO study. The genotyping in the NEO study was supported by the Centre National de Génotypage (Paris, France), headed by Jean-Francois Deleuze.

NFBC 1966 and 1986 We thank the late professor Paula Rantakallio (launch of NFBC1966), the participants in the 31- and 46-years-old study and the NFBC project center. We thank professor Anna-Liisa Hartikainen (launch of NFBC1986), the participants in the study and the NFBC project center.

NTR We would like to thank all twins and their families whom are cooperating in our research. We would like to thank the Avera Human Genetics research lab for their close collaboration, DNA typing and sample logistics.

ORCADES DNA extractions were performed at the Wellcome Trust Clinical Research Facility in Edinburgh. We would like to acknowledge the invaluable contributions of Lorraine Anderson and the research nurses in Orkney, the administrative team in Edinburgh and the people of Orkney.

PANIC We are grateful to the children and their parents for participating in the PANIC study and to the whole research team for their contribution in carrying out the study.

RAINE authors are grateful to the Raine Study participants and their families, and to the Raine Study research staff for cohort coordination and data collection. The authors gratefully acknowledge the

NH&MRC for their long term contribution to funding the study over the last 25 years and also the following Institutions for providing funding for Core Management of the Raine Study: The University of Western Australia (UWA), Raine Medical Research Foundation, UWA Faculty of Medicine, Dentistry and Health Sciences, Telethon Kids Institute and Women and Infants Research Foundation (King Edward Memorial Hospital), Curtin University and Edith Cowan University. The authors gratefully acknowledge the assistance of the Western Australian DNA Bank (National Health and Medical Research Council of Australia National Enabling Facility). This work was supported by resources provided by the Pawsey Supercomputing Centre with funding from the Australian Government and the Government of Western Australia.

TDCOB The Danish Childhood Obesity Biobank would like to thank all children, youths, and their families for participation in the studies and thus providing opportunities to detect novel insights in regards to childhood obesity.

TEENAGE We would like to thank all study participants and their families as well as all volunteers for their contribution in this study. We thank the following staff from the Sample Management and Genotyping Facilities at the Wellcome Trust Sanger Institute for sample preparation, quality control and genotyping: Dave Jones, Doug Simpkin, Emma Gray, Hannah Blackburn and Sarah Edkins.

YFS gratefully acknowledge the expert technical assistance in the statistical analyses by Ville Aalto and Irina Lisinen.

CHARGE Consortium Hematology Working Group Details of the acknowledgements are described in Ganesh et al, *Nat Genet* (2009)⁸⁰.

UK Biobank This study has been conducted using the UK Biobank Resource.